BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-(Mouse) Nr4a2 Antibody (Center)

Catalog Number: M01592

About Nr4a2

This enzyme scavenge exogenous and endogenous cytidine and 2'-deoxycytidine for UMP synthesis.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-(Mouse) Nr4a2 Antibody (Center)
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-(Mouse) Nr4a2 Antibody (Center) (Catalog # M01592). Tested in WB, IHC-P application(s). This antibody reacts with Mouse, Rat.
Application	IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q06219

Technical Details


Immunogen	This mouse Nr4a2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 164-197 amino acids from the Central region of mouse Nr4a2.
Predicted Reactive Species	Bovine, Chicken, Horse, Monkey, Pig, Sheep, Xenopus
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:2000 IHC-P: 1:25

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


	Tested Species: In-house tested species with positive results. Other applications have not been tested. Optimal dilutions should be determined by end users.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Anti-(Mouse) Nr4a2 Antibody (Center) (M01592) Images




All lanes : Anti-Nr4a2 Antibody (Center) at 1:2000 dilutionLane 1: mouse brain lysatesLane 2: rat brain lysatesLysates/proteins at 20 µg per lane. SecondaryGoat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilutionPredicted band size : 67 kDaBlocking/Dilution buffer: 5% NFDM/TBST.



Anti-Nr4a2 Antibody (Center)at 1:2000 dilution + mouse pancreas lysatesLysates/proteins at 20 µg per lane. SecondaryGoat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilutionPredicted band size : 67 kDaBlocking/Dilution buffer: 5% NFDM/TBST.



M01592 staining (Mouse) Nr4a2 in mouse brain sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0. 5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



M01592 staining (Mouse) Nr4a2 in mouse spleen sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0. 5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

For Research Use Only. Not for use in diagnostic procedures.