

Anti-IDO1/Ido Monoclonal Antibody

Catalog Number: M01705-1

About IDO1

Indoleamine 2, 3-dioxygenase1 (IDO1) is a 41-42 kD intracellular enzyme that catabolizes tryptophan into kynurenine. IDO1 modulates levels of the amino acid tryptophan, which is vital for cell growth, but is also involved in the suppression of the immune response. IDO1 effects on immune suppression are due to decreased tryptophan availability and the generation of tryptophan metabolites, resulting in negative effects on T lymphocytes, including proliferation, function and survival. IDO1 may be involved in the suppression of the immune response to tumors, and blocking the IDO1 pathway may be a potential target for immuno and cancer therapy. IDO1 is expressed in a wide variety of tissues and can be upregulated by interferon gamma and other inflammatory cytokines.

Overview


Product Name	Anti-IDO1/Ido Monoclonal Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-IDO1/Ido Monoclonal Antibody (Catalog # M01705-1). Tested in ELISA, WB applications. This antibody reacts with Human, Mouse.
Application	ELISA, IP, IF, IHC, WB
Clonality	Monoclonal Clone: 10.1 IgG3
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship on dry ice.)
Host	Mouse
Uniprot ID	P14902

Technical Details

Immunogen	Anti-IDO1 (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with fragment of recombinant human and mouse IDO1 protein followed by hybridoma development.
Predicted Reactive Species	Chimpanzee
Isotype	IgG3
Form	Liquid (sterile filtered)
Concentration	1.09 mg/mL by UV absorbance at 280 nm

Purification	Anti-IDO1 was purified from concentrated tissue culture supernate by Protein G chromatography followed by extensive dialysis against the buffer stated above. This antibody is specific for human and mouse IDO1 protein. A BLAST analysis was used to suggest cross-reactivity with IDO1 from human and mouse sources based on 100% homology with the immunizing sequence. Cross-reactivity with IDO1 from other sources has not been determined.
Suggested Dilutions	ELISA: 1:5000-1:50000 IHC: 1:100-1:500 IF Microscopy: 1:50-1:100 IP: 10-100 μ L WB: 1:500-1:1500 Anti-IDO1 antibody has been tested in ELISA, IP, and Western Blot. This antibody is suitable for use in IHC and Flow Cytometry. Specific conditions for reactivity should be optimized by the end user.

Anti-IDO1/Ido Monoclonal Antibody (M01705-1) Images



Western Blot of Mouse Anti-IDO1 Antibody. Lane 1: untreated HeLa cells . Lane 2: IFN-r treated HeLa cells. Load: 35 µg per lane. Primary antibody: IDO 1 Antibody at 1:1000 for overnight at 4°C. Secondary antibody: IRDye800™ mouse secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 41-42 kDa, 41-42 kDa for IDO-1. Other band(s): none.

2 Publications Citing This Product

1. PubMed ID: 27418932, Umbilical Cord Tissue-Derived Mesenchymal Stem Cells Induce T Lymphocyte Apoptosis and Cell Cycle Arrest by Expression of Indoleamine 2, 3-Dioxygenase

2. PubMed ID: 19948041, Expression of indoleamine 2, 3-dioxygenase in nasopharyngeal carcinoma impairs the cytolytic function of peripheral blood lymphocytes

Visit bosterbio.com/anti-ido1-antibody-m01705-1-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IDO1/Ido Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.