

Anti-HGD Antibody Picoband® (monoclonal, 2F11E1) APC Conjugated

Catalog Number: M01909-1-APC

About HGD

The HGD gene encodes homogentisate 1,2-dioxygenase (HGD), an enzyme involved in the catabolism of phenylalanine and tyrosine. This enzyme is involved in the catabolism of the amino acids tyrosine and phenylalanine. Mutations in this gene are the cause of the autosomal recessive metabolism disorder alkaptonuria. This gene is mapped to chromosome 3q21-q23 by a preliminary PCR screen of hamster/human somatic cell hybrid genomic DNA samples and by fluorescence in situ hybridization.

Overview

Product Name	Anti-HGD Antibody Picoband® (monoclonal, 2F11E1) APC Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (Flow Cytometry, IHC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Monoclonal 2F11E1
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Mouse
Uniprot ID	Q93099

Technical Details

Immunogen	E. coli-derived human HGD recombinant protein (Position: D374-N445).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	APC Excitation Wavelength: 633-647 nm Emission Wavelength: 660 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HGD Antibody (monoclonal, 2F11E1) - APC

For Research Use Only. Not for use in diagnostic procedures.