

Anti-Calpain 1 Antibody Picoband® (monoclonal, 2I10) APC Conjugated

Catalog Number: M01943-3-APC

About CAPN1

CAPN1 is also known as CANP or muCL. The calpains, calcium-activated neutral proteases, are nonlysosomal, intracellular cysteine proteases. The mammalian calpains include ubiquitous, stomach-specific, and muscle-specific proteins. The ubiquitous enzymes consist of heterodimers with distinct large, catalytic subunits associated with a common small, regulatory subunit. This gene encodes the large subunit of the ubiquitous enzyme, calpain 1. Several transcript variants encoding two different isoforms have been found for this gene.

Overview

Product Name	Anti-Calpain 1 Antibody Picoband® (monoclonal, 2I10) APC Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (Flow Cytometry, IF, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Monoclonal 2I10
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Mouse
Uniprot ID	P07384

Technical Details

Immunogen	E.coli-derived human Calpain 1 recombinant protein (Position: Q396-A555). Human Calpain 1 shares 86% amino acid (aa) sequence identity with both mouse and rat Calpain 1.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2a
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	APC Excitation Wavelength: 633-647 nm Emission Wavelength: 660 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Calpain 1 Antibody (monoclonal, 2I10) - APC

For Research Use Only. Not for use in diagnostic procedures.