

## Anti-Hu CD7 Purified Monoclonal Antibody

Catalog Number: M01974-1

### About CD7

CD7, also known as gp40, is a member of the immunoglobulin superfamily found on T cells, NK cells, thymocytes, hematopoietic progenitors, and monocytes (weakly). CD7 is also expressed on acute lymphocytic leukemia (ALL). CD7 crosslinking induces a calcium flux in T lymphocytes, presumably as a result of cytoplasmic domain association with PI3-kinase. CD7 co-stimulation can induce cytokine secretion and modulate cellular adhesion. A ligand of CD7, epithelial cell secreted protein K12, is produced in thymus to regulate thymocyte signaling and cytokine release. In lung microvascular endothelial cells CD7 serves as an IgM Fc receptor. Expression of CD7 is an important marker used in leukemia diagnostics.

### Overview

Product Name	Anti-Hu CD7 Purified Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Hu CD7 Purified Monoclonal Antibody (Catalog# M01974-1). Tested in Flow Cytometry, IP, WB, IHC-P application(s). This antibody reacts with Human.
Application	Flow Cytometry, IP, IHC-P, WB
Clonality	Monoclonal MEM-186
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage Instructions	Store at 2-8°C. Do not freeze.
Host	Mouse
Uniprot ID	P09564


### Technical Details

Immunogen	Human acute myelogenous leukaemia cell line KG-1. The antibody MEM-186 reacts with an extracellular epitope of CD7, a 40 kD type I transmembrane glycoprotein expressed on peripheral blood T lymphocytes, NK-cells, hematopoietic progenitors, monocytes (weakly) and also on acute lymphocytic leukemia.
Predicted Reactive Species	Primate
Cross Reactivity	This antibody does not cross-react with Thy-1.1 alloantigen.
Isotype	Mouse IgG1
Form	Liquid
Concentration	1 mg/ml
Purification	Purified by protein-A affinity chromatography.


Suggested Dilutions

Immunohistochemistry (paraffin sections): 5 ug/ml; positive tissue: spleen.  
Flow cytometry: 2 ug/ml.  
Western blotting: Non-reducing conditions.

## Anti-Hu CD7 Purified Monoclonal Antibody (M01974-1) Images



Separation of human CD7 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD7 (MEM-186) purified antibody (concentration in sample 0,33 µg/ml, GAM APC).



Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD7 (MEM-186) purified antibody (concentration in sample 0,33 µg/ml, GAM APC).

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Hu CD7 Purified Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.