

## Anti-RPA32/RPA2 Antibody Picoband® (monoclonal, 3B2E9) HRP Conjugated

Catalog Number: M02067-2-HRP

### About RPA2

This gene encodes a subunit of the heterotrimeric Replication Protein A (RPA) complex, which binds to single-stranded DNA (ssDNA), forming a nucleoprotein complex that plays an important role in DNA metabolism, being involved in DNA replication, repair, recombination, telomere maintenance, and co-ordinating the cellular response to DNA damage through activation of the ataxia telangiectasia and Rad3-related protein (ATR) kinase. The RPA complex protects single-stranded DNA from nucleases, prevents formation of secondary structures that would interfere with repair, and co-ordinates the recruitment and departure of different genome maintenance factors. The heterotrimeric complex has two different modes of ssDNA binding, a low-affinity and high-affinity mode, determined by which oligonucleotide/oligosaccharide-binding (OB) domains of the complex are utilized, and differing in the length of DNA bound. This subunit contains a single OB domain that participates in high-affinity DNA binding and also contains a winged helix domain at its carboxy terminus, which interacts with many genome maintenance protein. Post-translational modifications of the RPA complex also plays a role in co-ordinating different damage response pathways.

### Overview

Product Name	Anti-RPA32/RPA2 Antibody Picoband® (monoclonal, 3B2E9) HRP Conjugated
Reactive Species	Human
Application	WB, IHC, ELISA
Clonality	Monoclonal 3B2E9
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing.
Host	Mouse
Uniprot ID	P15927

### Technical Details

Immunogen	E.coli-derived human RPA32/RPA2 recombinant protein (Position: Q34-H254).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2a
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	HRP
Suggested Dilutions	Western blot, Optimal dilutions should be determined by end users. Immunohistochemistry (Paraffin-embedded Section), Optimal dilutions should be determined by end users. ELISA, Optimal dilutions should be determined by end users.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-RPA32/RPA2 Antibody (monoclonal, 3B2E9) - HRP

For Research Use Only. Not for use in diagnostic procedures.