

## Anti-Junctional Adhesion Molecule 1 (F11R) Mouse Monoclonal Antibody [Clone ID: OTI3H3]

Catalog Number: M02068-2


### Overview

Product Name	Anti-Junctional Adhesion Molecule 1 (F11R) Mouse Monoclonal Antibody [Clone ID: OTI3H3]
Reactive Species	Human
Description	Boster Bio F11R mouse monoclonal antibody, clone OTI3H3 (formerly 3H3). Catalog# M02068-2. Tested in IF, IHC, WB. This antibody reacts with Human.
Conjugate	Unconjugated
Application	IF, IHC, WB
Clonality	Monoclonal OTI3H3
Formulation	PBS (pH 7.3) containing 1% stabilizing protein, 50% glycerol and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q9Y624


### Technical Details

Immunogen	Full length human recombinant protein of human F11R (NP_653087) produced in HEK293T cell.
Isotype	IgG2b
Concentration	1 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	WB: 1:4000 IHC: 1:150 IF: 1:100


## Anti-Junctional Adhesion Molecule 1 (F11R) Mouse Monoclonal Antibody [Clone ID: OTI3H3] (M02068-2) Images




Anti-F11R mouse monoclonal antibody (M02068-2) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY F11R.



Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-F11R mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-F11R mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY F11R (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-F11R.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



For Research Use Only. Not for use in diagnostic procedures.