

Anti-Hu CD42b Purified GP1BA Monoclonal Antibody

Catalog Number: M02073

About GP1BA

CD42b (GPIb alpha) composes together with GPIb beta, GPIX and GPV the GPIb-IX-V receptor complex critical in the process of platelet-rich thrombus formation by tethering the platelet to a thrombogenic surface. CD42b binds to von Willebrand factor (VWF) exposed at a site of vascular injury, as well as to thrombin, coagulation factors XI and XII, high molecular weight kininogen, TSP-1, integrin Mac-1 and P-selectin. The extracellular domain of CD42b by its interactions also contributes to metastasis.

Overview

Product Name	Anti-Hu CD42b Purified GP1BA Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Hu CD42b Purified GP1BA Monoclonal Antibody (Catalog# M02073). Tested in Flow Cytometry, IP, WB, ICC, ELISA application(s). This antibody reacts with Human.
Application	ELISA, Flow Cytometry, IP, ICC, WB
Clonality	Monoclonal AK2
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage Instructions	Store at 2-8°C. Do not freeze.
Host	Mouse
Uniprot ID	P07359


Technical Details

Immunogen	Human platelets. The mouse monoclonal antibody AK2 recognizes an extracellular epitope of CD42b (GPIb alpha), a 135-145 kDa membrane glycoprotein expressed on platelets and megakaryocytes. CD42b and CD42c (GPIb beta) are composed in a disulfide linked heterodimer (CD42b/c; 160 kDa); CD42b/c forms a noncovalent complex with CD42a and CD42d.
Predicted Reactive Species	Primate
Cross Reactivity	This antibody does not cross-react with Thy-1.1 alloantigen.
Isotype	Mouse IgG1
Form	Liquid
Concentration	1 mg/ml
Purification	Purified by protein-A affinity chromatography.


Suggested Dilutions

Flow cytometry: 1-4 ug/ml.

Anti-Hu CD42b Purified GP1BA Monoclonal Antibody (M02073) Images



Separation of human thrombocytes (red-filled) from CD42b negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD42b (AK2) purified antibody (concentration in sample 4 µg/ml) GAM APC.



Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD42b (AK2) purified antibody (concentration in sample 4 µg/ml) GAM APC.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Hu CD42b Purified GP1BA Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.