

## Anti-HIF1AN Antibody (C-term)

Catalog Number: M02116-1

### About HIF1AN

Hydroxylates HIF-1 alpha at 'Asp-803' in the C-terminal transactivation domain (CAD). Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300- interacting transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD) of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively regulates NOTCH1 activity, accelerating myogenic differentiation. Positively regulates ASB4 activity, promoting vascular differentiation.

### Overview


Product Name	Anti-HIF1AN Antibody (C-term)
Reactive Species	Human
Description	Boster Bio Anti-HIF1AN Antibody (C-term) (Catalog # M02116-1). Tested in WB, IF application(s). This antibody reacts with Human.
Application	IF, WB
Clonality	Monoclonal 931CT3.1.1
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q9NWT6

### Technical Details


Immunogen	This HIF1AN antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 1-349 amino acids from human HIF1AN.
Predicted Reactive Species	Bovine
Isotype	IgG1,k
Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	IF: 1:25 WB: 1:4000



## Anti-HIF1AN Antibody (C-term) (M02116-1) Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling HIF1AN with M02116-1 at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-mouse IgG secondary antibody at 1/200 dilution (green). Immunofluorescence image showing mitochondrion staining on HeLa cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin at 1/100 dilution (red). The nuclear counter stain is DAPI (blue).



All lanes : Anti-HIF1AN Antibody (C-term) at 1:4000 dilution  
Lane 1: 293 whole cell lysate  
Lane 2: Jurkat whole cell lysate  
Lane 3: ZR-75-1 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 40 kDa  
Blocking/Dilution buffer: 5% NFDN/TBST.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HIF1AN Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.