

Anti-ADK Antibody Picoband® (monoclonal, 7F4) PE Conjugated

Catalog Number: M02193-PE

About ADK

This gene is an enzyme which catalyzes the transfer of the gamma-phosphate from ATP to adenosine, thereby serving as a regulator of concentrations of both extracellular adenosine and intracellular adenosine nucleotides. Adenosine has widespread effects on the cardiovascular, nervous, respiratory, and immune systems and inhibitors of the enzyme could play an important pharmacological role in increasing intravascular adenosine concentrations and acting as anti-inflammatory agents. Multiple transcript variants encoding different isoforms have been found for this gene.

Overview

Product Name	Anti-ADK Antibody Picoband® (monoclonal, 7F4) PE Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (Flow Cytometry, IHC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Monoclonal 7F4
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Mouse
Uniprot ID	P55263

Technical Details

Immunogen	E. coli-derived human ADK recombinant protein (Position: K165-T351). Human ADK shares 88.8% and 88.2% amino acid (aa) sequence identity with mouse and rat ADK, respectively.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG1
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	PE Excitation Wavelength: 566 nm Emission Wavelength: 574 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ADK Antibody (monoclonal, 7F4) - PE

For Research Use Only. Not for use in diagnostic procedures.