

Anti-IFITM3 Antibody (N-term)

Catalog Number: M02265-1

About IFITM3

IFN-induced antiviral protein which disrupts intracellular cholesterol homeostasis. Inhibits the entry of viruses to the host cell cytoplasm by preventing viral fusion with cholesterol depleted endosomes. May inactivate new enveloped viruses which buds out of the infected cell, by letting them go out with a cholesterol depleted membrane. Active against multiple viruses, including influenza A virus, SARS coronavirus (SARS-CoV), Marburg virus (MARV) and Ebola virus (EBOV), Dengue virus (DNV), West Nile virus (WNV), human immunodeficiency virus type 1 (HIV-1) and vesicular stomatitis virus (VSV). Can inhibit: influenza virus hemagglutinin protein-mediated viral entry, MARV and EBOV GP1,2- mediated viral entry, SARS-CoV S protein-mediated viral entry and VSV G protein-mediated viral entry. Plays a critical role in the structural stability and function of vacuolar ATPase (v-ATPase). Establishes physical contact with the v-ATPase of endosomes which is critical for proper clathrin localization and is also required for the function of the v-ATPase to lower the pH in phagocytic endosomes thus establishing an antiviral state.


Overview

Product Name	Anti-IFITM3 Antibody (N-term)
Reactive Species	Human
Description	Boster Bio Anti-IFITM3 Antibody (N-term) (Catalog # M02265-1). Tested in WB application(s). This antibody reacts with Human.
Application	WB
Clonality	Monoclonal 1082CT14.1.3.1
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q01628

Technical Details

Immunogen	This IFITM3 antibody is generated from a mouse immunized with a recombinant protein KLH conjugated synthetic peptide between 1-133 amino acids from the N-terminal region of human IFITM3.
Predicted Reactive Species	Human
Isotype	IgG1,k
Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	WB: 1:4000

Anti-IFITM3 Antibody (N-term) (M02265-1) Images



All lanes : Anti-Fragilis (IFITM3) Antibody (N-term) at 1:4000 dilution
Lane 1: NCI-H460 whole cell lysate
Lane 2: HepG2 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 15 kDa
Blocking/Dilution buffer: 5% NFD/MTBST.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IFITM3 Antibody (N-term)

For Research Use Only. Not for use in diagnostic procedures.