

Anti-HSPB8/Hsp22 Antibody Picoband™ (monoclonal, 7D8)

Catalog Number: M02492-2

About HSPB8

Heat shock protein beta-8 is a protein that in humans is encoded by the HSPB8 gene. The protein encoded by this gene belongs to the superfamily of small heat-shock proteins containing a conservative alpha-crystallin domain at the C-terminal part of the molecule. The expression of this gene is induced by estrogen in estrogen receptor-positive breast cancer cells, and this protein also functions as a chaperone in association with Bag3, a stimulator of macroautophagy. Thus, this gene appears to be involved in regulation of cell proliferation, apoptosis, and carcinogenesis, and mutations in this gene have been associated with different neuromuscular diseases, including Charcot-Marie-Tooth disease.

Overview

Product Name	Anti-HSPB8/Hsp22 Antibody Picoband™ (monoclonal, 7D8)
Reactive Species	Human, Rat
Description	Boster Bio Anti-HSPB8/Hsp22 Antibody Picoband™ (monoclonal, 7D8) catalog # M02492-2. Tested in Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human, Rat.
Application	Flow Cytometry, IF, ICC, WB
Clonality	Monoclonal 7D8
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q9UJY1

Technical Details

Immunogen	E.coli-derived human HSPB8/Hsp22 recombinant protein (Position: M1-T196). Human HSPB8/Hsp22 shares 94.4% and 95.4% amino acid (aa) sequence identity with mouse and rat HSPB8/Hsp22, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human, Rat</p> <p>Immunocytochemistry/Immunofluorescence, 5ug/ml, Human</p> <p>Flow Cytometry, 1-3ug/1x10⁶ cells, Human</p>

Anti-HSPB8/Hsp22 Antibody Picoband™ (monoclonal, 7D8) (M02492-2) Images




Figure 1. Western blot analysis of HSPB8/Hsp22 using anti-HSPB8/Hsp22 antibody (M02492-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.
Lane 1: human HeLa whole cell lysates;
Lane 2: human T-47D whole cell lysates;
Lane 3: rat RH35 whole cell lysates.
After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-HSPB8/Hsp22 antigen affinity purified monoclonal antibody (Catalog # M02492-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for HSPB8/Hsp22 at approximately 22KD. The expected band size for HSPB8/Hsp22 is at 22KD.




Figure 2. Flow Cytometry analysis of CACO-2 cells using anti-HSPB8/Hsp22 antibody (M02492-2). Overlay histogram showing CACO-2 cells stained with M02492-2 (Blue line).The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-HSPB8/Hsp22 Antibody (M02492-2, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.





Figure 3. Flow Cytometry analysis of U2OS cells using anti-HSPB8/Hsp22 antibody (M02492-2). Overlay histogram showing U2OS cells stained with M02492-2 (Blue line).The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-HSPB8/Hsp22 Antibody (M02492-2, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Figure 4. IF analysis of HSPB8/Hsp22 using anti-HSPB8/Hsp22 antibody (M02492-2).



HSPB8/Hsp22 was detected in immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5ug/mL mouse anti-HSPB8/Hsp22 Antibody (M02492-2) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HSPB8/Hsp22 Antibody (monoclonal, 7D8)