

Anti-CoREST (RCOR1) Mouse Monoclonal Antibody [Clone ID: OTI9F6]

Catalog Number: M02799


Overview

Product Name	Anti-CoREST (RCOR1) Mouse Monoclonal Antibody [Clone ID: OTI9F6]
Reactive Species	Human, Mouse, Rat
Description	Boster Bio RCOR1 mouse monoclonal antibody, clone OTI9F6. Catalog# M02799. Tested in IHC, WB. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Monoclonal OTI9F6
Formulation	PBS (pH 7.3) containing 1% stabilizing protein, 50% glycerol and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C as received.
Host	Mouse
Uniprot ID	Q9UKL0


Technical Details

Immunogen	Human recombinant protein fragment corresponding to amino acids 256-485 of human RCOR1 (NP_055971) produced in E.coli.
Isotype	IgG1
Concentration	1 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	WB: 1:500 IHC: 1:500


Anti-CoREST (RCOR1) Mouse Monoclonal Antibody [Clone ID: OT19F6] (M02799) Images




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RCOR1 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RCOR1 (1:500).



Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-RCOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min)



Immunohistochemical staining of paraffin-embedded Human appendix tissue within the normal limits using anti-RCOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min)



Immunohistochemical staining of paraffin-embedded Human lung tissue within the normal limits using anti-RCOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min)

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



For Research Use Only. Not for use in diagnostic procedures.