BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-TRRAP Antibody (C-term)

Catalog Number: M02981

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview


Product Name	Anti-TRRAP Antibody (C-term)
Reactive Species	Human
Description	Boster Bio Anti-TRRAP Antibody (C-term) (Catalog # M02981). Tested in WB, IF application(s). This antibody reacts with Human.
Application	IF, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9Y4A5

Technical Details


Immunogen	This TRRAP antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 3015-3049 amino acids from the C-terminal region of human TRRAP.
Predicted Reactive Species	Bovine, Mouse
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	IF: 1:25 WB: 1:2000

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Anti-TRRAP Antibody (C-term) (M02981) Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling TRRAP with M02981 at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG secondary antibody at 1/200 dilution (green). Immunofluorescence image showing nucleus staining on HeLa cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin at 1/100 dilution (red).



All lanes : Anti-TRRAP Antibody (C-term) at 1:2000 dilutionLane 1: K562 whole cell lysatesLane 2: Hela whole cell lysatesLysates/proteins at 20 µg per lane. SecondaryGoat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 438 kDaBlocking/Dilution buffer: 5% NFDM/TBST.



Anti-TRRAP Antibody (C-term)at 1:2000 dilution + K562 whole cell lysatesLysates/proteins at 20 µg per lane. SecondaryGoat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 438 kDaBlocking/Dilution buffer: 5% NFDM/TBST.

Anti-TRRAP Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.