

## Anti-DYNC1H1 Antibody (C-term)

Catalog Number: M03059

### About DYNC1H1

Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. Dynein has ATPase activity; the force-producing power stroke is thought to occur on release of ADP.


### Overview

Product Name	Anti-DYNC1H1 Antibody (C-term)
Reactive Species	Human, Mouse
Description	Boster Bio Anti-DYNC1H1 Antibody (C-term) (Catalog # M03059). Tested in WB, IHC-P application(s). This antibody reacts with Human, Mouse.
Application	IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q14204


### Technical Details

Immunogen	This DYNC1H1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 4202-4236 amino acids from the C-terminal region of human DYNC1H1.
Predicted Reactive Species	Bovine, Chicken, Horse, Monkey, Pig, Sheep, Xenopus
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:2000 IHC-P: 1:25


## Anti-DYNC1H1 Antibody (C-term) (M03059) Images



All lanes : Anti-DYNC1H1 Antibody (C-term) at 1:2000 dilution  
Lane 1: U-87 MG whole cell lysates  
Lane 2: PC-3 whole cell lysates  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution  
Predicted band size : 532 kDa  
Blocking/Dilution buffer: 5% NFD/MTBST.



All lanes : Anti-DYNC1H1 Antibody (C-term) at 1:2000 dilution  
Lane 1: A431 whole cell lysate  
Lane 2: PC-3 whole cell lysate  
Lane 3: mouse brain lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 532 kDa  
Blocking/Dilution buffer: 5% NFD/MTBST.



M03059 staining DYNC1H1 in human testis sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

### Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-DYNC1H1 Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.