

Anti-Aconitase 2 (ACO2) Mouse Monoclonal Antibody [Clone ID: OTI3G8]

Catalog Number: M03096

Overview


Product Name	Anti-Aconitase 2 (ACO2) Mouse Monoclonal Antibody [Clone ID: OTI3G8]
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ACO2 (Aconitase 2) mouse monoclonal antibody, clone OTI3G8 (formerly 3G8). Catalog# M03096. Tested in IF, IHC, WB. This antibody reacts with Human, Mouse, Rat.
Conjugate	Unconjugated
Application	IF, IHC, WB
Clonality	Monoclonal OTI3G8
Formulation	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q99798

Technical Details


Immunogen	Full length human recombinant protein of human ACO2 (NP_001089) produced in HEK293T cell.
Isotype	IgG1
Concentration	0.9 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows:




Anti-Aconitase 2 (ACO2) Mouse Monoclonal Antibody [Clone ID: OTI3G8] (M03096) Images




Anti-ACO2 mouse monoclonal antibody (M03096) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ACO2.




Immunofluorescent staining of HepG2 cells using anti-ACO2 mouse monoclonal antibody (M03096).



Immunohistochemical staining of paraffin-embedded Ovary tissue within the normal limits using anti-ACO2mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of breast tissue using anti-ACO2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer




Immunohistochemical staining of paraffin-embedded Kidney tissue within the normal limits using anti-ACO2mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ACO2 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACO2.



Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-ACO2 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Aconitase 2 (ACO2) Mouse Monoclonal Antibody [Clone ID: OTI3G8]