

Anti-Fibrillarin FBL Rabbit Monoclonal Antibody

Catalog Number: M03178

About FBL

Receptor for bradykinin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system.

Overview

Product Name	Anti-Fibrillarin FBL Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Fibrillarin FBL Rabbit Monoclonal Antibody catalog # M03178. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Conjugate	FITC
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal DIO-6
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P22087

Technical Details

Immunogen	A synthesized peptide derived from human Fibrillarin
Predicted Reactive Species	Human, Primate
Cross Reactivity	Detects ~20kDa. Does not cross-react with alphaB-crystallin, betaL-crystallin, ̢H-crystallin, gamma-crystallin, HSP25, HSP27 or HSP47 proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

WB 1:500-1:2000

IHC 1:50-1:100


ICC/IF 1:50-1:100

IP:


1:30

FC 1:50


Anti-Fibrillarin FBL Rabbit Monoclonal Antibody (M03178) Images




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot analysis of Fibrillarin expression in HepG2 cell lysate.





All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Immunohistochemical analysis of paraffin-embedded human testis, using Fibrillarin Antibody.

Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis of HeLa cells, using Fibrillarin Antibody .

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Fibrillarin FBL Rabbit Monoclonal Antibody