

Anti-Glutamine Synthetase (GLUL) Mouse Monoclonal Antibody [Clone ID: OTI1F4]

Catalog Number: M03191-1


Overview

Product Name	Anti-Glutamine Synthetase (GLUL) Mouse Monoclonal Antibody [Clone ID: OTI1F4]
Reactive Species	Human, Mouse, Rat
Description	Boster Bio GLUL mouse monoclonal antibody, clone OTI1F4 (formerly 1F4). Catalog# M03191-1. Tested in FC, IF, IHC, WB. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IF, IHC, WB
Clonality	Monoclonal OTI1F4
Formulation	PBS (pH 7.3) containing 1% stabilizing protein, 50% glycerol and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C as received.
Host	Mouse
Uniprot ID	P15104


Technical Details

Immunogen	Full length human recombinant protein of human GLUL (NP_002056) produced in HEK293T cell.
Isotype	IgG2a
Concentration	1 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	WB: 1:1000~2000 IHC: 1:50 IF: 1:50~100 Flow cytometry: 1:100


Anti-Glutamine Synthetase (GLUL) Mouse Monoclonal Antibody [Clone ID: OTI1F4] (M03191-1) Images




Western blot analysis of GLUL using anti-GLUL antibody (M03191-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human K562 whole cell lysates, Lane 2: rat brain tissue lysates, Lane 3: rat liver tissue lysates, Lane 4: mouse brain tissue lysates, Lane 5: mouse liver tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-GLUL antigen affinity purified monoclonal antibody (Catalog # M03191-1) at at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for GLUL at approximately 42 kDa. The expected band size for GLUL is at 42 kDa.




Immunofluorescent staining of HepG2 cells using anti-GLUL mouse monoclonal antibody (M03191-1).




Flow cytometric Analysis of Jurkat cells




Anti-GLUL mouse monoclonal antibody (M03191-1) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY GLUL.




Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-GLUL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Flow cytometric Analysis of HeLa cells



Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-GLUL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)





Immunohistochemical staining of paraffin-embedded Human Ovary tissue within the normal limits using anti-GLUL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-GLUL monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).

Figure from citation: Western blot analysis of GLUL protein level by using anti-GLUL antibody in LCC9 cells were treated with 10058-F4 (25 uM) or vehicle for 48 h or transfected with MYC or control siRNA for 48 h. Knockdown of MYC increased GLS/GAC levels and decreased GLUL levels.



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GLUL (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GLUL.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Glutamine Synthetase (GLUL) Mouse Monoclonal Antibody [Clone ID: OT11F4]

For Research Use Only. Not for use in diagnostic procedures.