

## Anti-Cyclophilin B Antibody

Catalog Number: M03229-3

### About PPIB

PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.


### Overview

Product Name	Anti-Cyclophilin B Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Cyclophilin B Antibody (Catalog # M03229-3). Tested in WB, Flow Cytometry, IHC-P application(s). This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P23284


### Technical Details

Immunogen	This Cyclophilin B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 161-195 amino acids from the human region of human Cyclophilin B.
Predicted Reactive Species	Mouse
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:2000 IHC-P: 1:25 FC: 1:25


## Anti-Cyclophilin B Antibody (M03229-3) Images




All lanes : Anti-Cyclophilin B Antibody at 1:2000 dilution  
Lane 1: HeLa whole cell lysate  
Lane 2: HepG2 whole cell lysate  
Lane 3: A431 whole cell lysate  
Lane 4: BA/F3 whole cell lysate  
Lane 5: NIH/3T3 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa  
Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-Cyclophilin B Antibody at 1:2000 dilution  
Lane 1: HeLa whole cell lysate  
Lane 2: HepG2 whole cell lysate  
Lane 3: A431 whole cell lysate  
Lane 4: BA/F3 whole cell lysate  
Lane 5: NIH/3T3 whole cell lysate  
Lane 6: PC-12 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa  
Blocking/Dilution buffer: 5% NFDm/TBST.



M03229-3 staining Cyclophilin B in human placenta tissue sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing HeLa cells stained with M03229-3 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (M03229-3, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1g/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



**Anti-Cyclophilin B Antibody**

For Research Use Only. Not for use in diagnostic procedures.