

# **Anti-PDL2 PDCD1LG2 Monoclonal Antibody [4E10]**

Catalog Number: M03295

#### **About PDCD1LG2**

Cell-mediated immune responses are initiated by T lymphocytes that are themselves stimulated by co gnate peptides bound to MHC molecules on antigen-presenting cells (APC). T-cell activation is generally self-limited as activated T cells express receptors such as PD-1 (also known as PDCD-1) that mediate inhibitory signals from the APC. PD-1 can bind two different but related ligands, PD-L1 and PD-L2, both of which are thought act as a negative regulator of T cell activation. However, it has been suggested that PD-L2 can act to stimulate an immunogenic response through and alternative receptor from PD-1.

#### Overview

Product Name	Anti-PDL2 PDCD1LG2 Monoclonal Antibody [4E10]	
Reactive Species	Human	
Description	Boster Bio Anti-PDL2 PDCD1LG2 Monoclonal Antibody [4E10] (Catalog # M03295). Tested in ELISA, WB, IHC-P, ICC, IF, Flow Cytometry applications. This antibody reacts with Human.	
Application	ELISA, Flow Cytometry, IF, IHC-P, ICC, WB	
Clonality	Monoclonal Clone: 4E10	
Formulation	PD-L2 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol.	
Storage Instructions	PD-L2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. Avoid repeated freeze-thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	
Host	Mouse	
Uniprot ID	P9BQ51	

#### **Technical Details**

Immunogen	PD-L2 antibody was raised against the extracellular domain of human PD-L2.
Predicted Reactive Species	Mouse, Rat
Cross Reactivity	beta-Actin antibody is human, mouse, rat, rabbit, chicken, zebrafish and drosophila reactive.
Isotype	lgG1
Form	Liquid
Concentration	1 mg/mL
Purification	PD-L2 Antibody is supplied as protein A purified IgG1.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this



# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

:

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:


Boster Bio's internal QC testing used:

PD-L2 antibody can be used for detection of PD-L2 by Western blot at 0.5 - 1 ug/mL. Antibody can also be used for immunohistochemistry starting at 2 ug/mL. For immunofluorescence start at 20 ug/mL.


Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples; Immunocytochemistry in human samples; Immunofluorescence in human samples and Flow Cytometry in mouse samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher.




### Anti-PDL2 PDCD1LG2 Monoclonal Antibody [4E10] (M03295) Images



Western blot analysis of PD-L2 in overexpressing HEK293 cells PD-L2 antibody at 0.5 and 1 ug/ml




Immunocytochemistry of PD-L2 in transfected HEK293 cells with PD-L2 antibody at 5 ug/mL. Lower left: Immunocytochemistry in transfected HEK293 cells with control mouse IgG antibody at 5 ug/mL.



Immunofluorescence of PD-L2 in transfected HEK293 cells with PD-L2 antibody at 20 ug/mL.

Green: PDL2 Antibody [4E10] (RF16021)

Blue: DAPI staining



Immunofluorescence of PD-L2 in human tonsil tissue with PD-L2 antibody at 20 ug/mL.

Green: PDL2 Antibody [4E10] (RF16021)


Blue: DAPI staining


Immunofluorescence of PD-L2 in human colon carcinoma tissue with PD-L2 antibody at 20 ug/mL.

Green: PDL2 Antibody [4E10] (M03295)


Blue: DAPI staining








Immunohistochemistry of PD-L2 in human tonsil tissue with PD-L2 antibody at 2 ug/mL.



Immunohistochemistry of PD-L2 in human colon carcinoma tissue with PD-L2 antibody at 2 ug/mL.



Flow cytometry analysis of PD-L2 overexpressing HEK293 cells using PD-L2 antibody and control mouse IgG antibody at 10 ug/ml. Blue: Untransfected HEK293 cells. Yellow: PD-L2 overexpressing HEK293 cells.

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.