

Anti-RNF20 Antibody

Catalog Number: M03457-2

About RNF20

Component of the RNF20/40 E3 ubiquitin-protein ligase complex that mediates monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1). H2BK120ub1 gives a specific tag for epigenetic transcriptional activation and is also prerequisite for histone H3 'Lys-4' and 'Lys-79' methylation (H3K4me and H3K79me, respectively). It thereby plays a central role in histone code and gene regulation. The RNF20/40 complex forms a H2B ubiquitin ligase complex in cooperation with the E2 enzyme UBE2A or UBE2B; reports about the cooperation with UBE2E1/UBCH are contradictory. Required for transcriptional activation of Hox genes. Recruited to the MDM2 promoter, probably by being recruited by p53/TP53, and thereby acts as a transcriptional coactivator. Mediates the polyubiquitination of isoform 2 of PA2G4 in cancer cells leading to its proteasome-mediated degradation.


Overview

Product Name	Anti-RNF20 Antibody
Reactive Species	Human
Description	Boster Bio Anti-RNF20 Antibody (Catalog # M03457-2). Tested in WB application(s). This antibody reacts with Human.
Application	WB
Clonality	Monoclonal 1594CT552.128.36
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q5VTR2

Technical Details

Immunogen	This RNF20 antibody is generated from a mouse immunized with a recombinant protein of human RNF20.
Predicted Reactive Species	Human
Isotype	IgG1,k
Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	WB: 1:2000

Anti-RNF20 Antibody (M03457-2) Images



All lanes : Anti-RNF20 Antibody at 1:2000 dilution
Lane 1: HeLa whole cell lysate
Lane 2: Jurkat whole cell lysate
Lane 3: HL-60 whole cell lysate
Lane 4: U87-MG whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 114 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-RNF20 Antibody

For Research Use Only. Not for use in diagnostic procedures.