

# Anti-Hu CD150 Purified Azide Free SLAMF1 Monoclonal Antibody

Catalog Number: M03988

#### **About SLAMF1**

CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail contains three unique tyrosine-based motifs. These binding sites can be recognized by SH2-binding phosphatases and the adaptor proteins, such as SAP/SH2D1A or EAT-2. The SLAM family receptors are involved in leucocyte activation and contribute to the effective germinal center formation, generation of high-affinity antibody-secreting plasma cells, and memory T and B cells, thereby facilitating long-term immune response. CD150 expression is upregulated after cell activation.

### Overview

Product Name	Anti-Hu CD150 Purified Azide Free SLAMF1 Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Hu CD150 Purified Azide Free SLAMF1 Monoclonal Antibody (Catalog# M03988). Tested in Flow Cytometry, IP, ICC application(s). This antibody reacts with Human.
Application	Flow Cytometry, IP, ICC
Clonality	Monoclonal SLAM.4
Formulation	Phosphate buffered saline (PBS), pH 7.4, azide-free
Storage Instructions	Store at 2-8°C. Do not freeze.
Host	Mouse
Uniprot ID	Q13291

### **Technical Details**

Immunogen	Human CD150-transfected 300.19 cells. The mouse monoclonal antibody SLAM.4 recognizes an extracellular epitope of CD150, a cell surface molecule expressed on lymphocytes and involved in their activation.
Predicted Reactive Species	Primate
Cross Reactivity	This antibody does not cross-react with Thy-1.1 alloantigen.
Isotype	Mouse IgG1
Form	Liquid
Concentration	1 mg/ml



888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Purified by protein-A affinity chromatography.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used: Flow cytometry: 1-5 ug/ml.

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Hu CD150 Purified Azide Free SLAMF1 Monoclonal Antibody