

## **Anti-SLAMF1** Rabbit Monoclonal Antibody

Catalog Number: M03988-1

#### **About SLAMF1**

Plays a major role in tight junction-specific obliteration of the intercellular space.

### Overview

| Product Name         | Anti-SLAMF1 Rabbit Monoclonal Antibody  |
|----------------------|---|
| Reactive Species     | Human   |
| Description          | Boster Bio Anti-SLAMF1 Rabbit Monoclonal Antibody catalog # M03988-1. Tested in WB, IP, Flow Cytometry applications. This antibody reacts with Human. |
| Conjugate            | FITC  |
| Application          | Flow Cytometry, IP, WB  |
| Clonality            | Monoclonal 24S06  |
| Formulation          | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.                                   |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.            |
| Host                 | Rabbit  |
| Uniprot ID           | Q13291  |

#### **Technical Details**

| Immunogen                  | A synthesized peptide derived from human SLAMF1  |
|----------------------------|--|
| Predicted Reactive Species | Human, Primate   |
| Cross Reactivity           | Detects ~20kDa. Does not cross-react with alphaB-crystallin, betaL-crystallin, ©H- crystallin, gamma-crystallin, HSP25, HSP27 or HSP47 proteins. |
| Isotype                    | IgG  |
| Form                       | Liquid   |
| Concentration              | Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.   |
| Purification               | Affinity-chromatography  |
| Suggested Dilutions        | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.                                      |




# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

| If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  WB 1:500-1:2000  IP 1:50  FC 1:50 |
|--|
|--|



### Anti-SLAMF1 Rabbit Monoclonal Antibody (M03988-1) Images



Western blot analysis of SLAMF1 expression in Jurkat cell lysate.

### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-SLAMF1 Rabbit Monoclonal Antibody