

Anti-Nectin (NDN) Mouse Monoclonal Antibody [Clone ID: OTI5D1]

Catalog Number: M04394


Overview

Product Name	Anti-Nectin (NDN) Mouse Monoclonal Antibody [Clone ID: OTI5D1]
Reactive Species	Human
Description	Boster Bio NDN mouse monoclonal antibody, clone OTI5D1 (formerly 5D1). Catalog# M04394. Tested in IF, IHC, WB. This antibody reacts with Human.
Application	IF, IHC, WB
Clonality	Monoclonal OTI5D1
Formulation	PBS (pH 7.3) containing 1% stabilizing protein, 50% glycerol and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C as received.
Host	Mouse
Uniprot ID	Q99608


Technical Details

Immunogen	Full length human recombinant protein of human NDN (NP_002478) produced in HEK293T cell.
Isotype	IgG2a
Concentration	1 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	WB 1:4000 IHC 1:150 IF 1:100


Anti-Necdin (NDN) Mouse Monoclonal Antibody [Clone ID: OTI5D1] (M04394) Images




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NDN (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NDN.




Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)


Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-NDN mouse




monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)


Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by



10mM citric buffer



Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-NDN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



Immunofluorescent staining of HeLa cells using anti-NDN mouse monoclonal antibody (M04394).

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Necdin (NDN) Mouse Monoclonal Antibody [Clone ID: OTI5D1]

For Research Use Only. Not for use in diagnostic procedures.