

Anti-Pirh2 RCHY1 Monoclonal Antibody

Catalog Number: M04533

About RCHY1

Anti Mesothelin Antibody recognizes Mesothelin that is a glycosyl-phosphatidylinositol-anchored glycoprotein present on the cell surface of various human solid tumors. The mesothelin (MSLN) gene encodes a 71-kDa precursor protein that is processed to a 40-kDa glycosylphosphatidylinositol-anchored protein that composes the mature portion and an NH2 terminal 31-kDa fragment called megakaryocyte-potentiating factor that is released from the cell. Mesothelin is a tumor differentiation antigen present at low levels on a restricted set of normal adult tissues, such as mesothelium, but aberrantly over expressed in mesotheliomas, ovarian, and pancreatic cancers. The biological functions of mesothelin remain elusive. A recent study showed that mesothelin binds to MUC16/CA125, and that this interaction mediates cell adhesion, suggesting that there may be an important role for MUC16/CA125 and mesothelin in the metastatic spread of ovarian cancer.

Overview

Product Name	Anti-Pirh2 RCHY1 Monoclonal Antibody
Reactive Species	Human, Rat
Description	Boster Bio Anti-Pirh2 RCHY1 Monoclonal Antibody catalog # M04533. Tested in ELISA, Flow Cytometry, IF, IHC, WB applications. This antibody reacts with Human, Rat.
Application	ELISA, Flow Cytometry, IF, IHC, WB
Clonality	Monoclonal 1H10
Formulation	Ascitic fluid containing 0.03% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q96PM5

Technical Details

Immunogen	Purified recombinant fragment of human Pirh2 expressed in E. Coli.
Predicted Reactive Species	Chimpanzee, Macaque
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml

Purification	Affinity purification
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>WB 1:500-1:2000</p> <p>IHC 1:200-1:1000</p> <p>IF 1:200-1:1000</p> <p>FC 1:200-1:400</p> <p>ELISA 1:10000</p>

Anti-Pirh2 RCHY1 Monoclonal Antibody (M04533) Images




Figure 4. Flow Cytometry validation of RCHY1 using Anti-Pirh2 RCHY1 Monoclonal Antibody (M04533).

Flow cytometric (FCM) analysis of PC-12 cells using Pirh2 Monoclonal Antibody (blue) and negative control (red).




Figure 3. Immunofluorescent staining data of RCHY1 using Anti-Pirh2 RCHY1 Monoclonal Antibody (M04533).

Immunofluorescence (IF) analysis of HeLa cells using Pirh2 Monoclonal Antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 2. Immunohistochemistry validation of RCHY1 using Anti-Pirh2 RCHY1 Monoclonal Antibody (M04533).

Immunohistochemistry (IHC) analysis of paraffin-embedded Human Tonsil tissues with AEC staining using Pirh2 Monoclonal Antibody.
For more protocol information of IHC




Figure 1. Western blotting validation for Anti-Pirh2 RCHY1 Monoclonal Antibody M04533

Western Blot (WB) analysis using Pirh2 Monoclonal Antibody against HeLa (1)

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Pirh2 RCHY1 Monoclonal Antibody