BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-RPS7 Antibody (C-Term)

Catalog Number: M04736

About RPS7

Required for rRNA maturation.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-RPS7 Antibody (C-Term)
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-RPS7 Antibody (C-Term) (Catalog # M04736). Tested in WB, Flow Cytometry, IF application(s). This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IF, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P62081

Technical Details


Immunogen	This RPS7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 158-191 amino acids from human RPS7.
Predicted Reactive Species	Bovine
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	IF: 1:25 WB: 1:2000

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


	FC: 1:25

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Anti-RPS7 Antibody (C-Term) (M04736) Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling RPS7 with M04736 at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG secondary antibody at 1/200 dilution (green). Immunofluorescence image mainly showing cytoplasm staining on HeLa cell line. The nuclear counter stain is DAPI (blue).



All lanes : Anti-RPS7 Antibody (C-Term) at 1:2000 dilutionLane 1: Hela whole cell lysateLane 2: A549 whole cell lysateLane 3: HepG2 whole cell lysateLane 4: Jurkat whole cell lysateLane 5: NIH/3T3 whole cell lysateLane 6: PC-12 whole cell lysateLysates/proteins at 20 µg per lane. SecondaryGoat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 22 kDaBlocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing Hela cells stained with M04736 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then icubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (M04736, 1:25 dilution) for 60 min at 37ºC. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at 37ºC. Isotype control antibody (blue line) was rabbit IgG (1g/1x10^6 cells) used under the same conditions. Acquisition of >10, 000 events was performed.

For Research Use Only. Not for use in diagnostic procedures.