

## Anti-RPS7 Antibody (C-Term)

Catalog Number: M04736

### About RPS7

Required for rRNA maturation.

### Overview


Product Name	Anti-RPS7 Antibody (C-Term)
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-RPS7 Antibody (C-Term) (Catalog # M04736). Tested in WB, Flow Cytometry, IF application(s). This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IF, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P62081

### Technical Details


Immunogen	This RPS7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 158-191 amino acids from human RPS7.
Predicted Reactive Species	Bovine
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>IF: 1:25 WB: 1:2000 FC: 1:25</p>



## Anti-RPS7 Antibody (C-Term) (M04736) Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling RPS7 with M04736 at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG secondary antibody at 1/200 dilution (green). Immunofluorescence image mainly showing cytoplasm staining on HeLa cell line. The nuclear counter stain is DAPI (blue).



All lanes : Anti-RPS7 Antibody (C-Term) at 1:2000 dilution

Lane 1: HeLa whole cell lysate

Lane 2: A549 whole cell lysate

Lane 3: HepG2 whole cell lysate

Lane 4: Jurkat whole cell lysate

Lane 5: NIH/3T3 whole cell lysate

Lane 6: PC-12 whole cell lysate


Lysates/proteins at 20 µg per lane.

Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.

Predicted band size : 22 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing HeLa cells stained with M04736 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (M04736, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1g/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-RPS7 Antibody (C-Term)