

## Anti-MAP3K4 Rabbit Monoclonal Antibody

Catalog Number: M04842


### Overview

|                      |  |
|----------------------|--|
| Product Name         | Anti-MAP3K4 Rabbit Monoclonal Antibody   |
| Reactive Species     | Human  |
| Description          | Boster Bio Anti-MAP3K4 Rabbit Monoclonal Antibody catalog # M04842. Tested in WB, IHC applications. This antibody reacts with Human.   |
| Application          | IHC, WB  |
| Clonality            | Monoclonal 28M41   |
| Formulation          | Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.<br>*This antibody is supplied in a stabilized formulation.<br>Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required. |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.   |
| Host                 | Rabbit   |
| Uniprot ID           | Q9Y6R4   |

### Technical Details

|                     |   |
|---------------------|---|
| Immunogen           | A synthesized peptide derived from human MAP3K4 |
| Isotype             | IgG   |
| Form                | Liquid  |
| Concentration       | 0.5mg/ml  |
| Purification        | Affinity-chromatography                         |
| Suggested Dilutions | WB 1:500-2000<br>IHC 1:50-200                   |

## Anti-MAP3K4 Rabbit Monoclonal Antibody (M04842) Images



Western blot analysis of MAP3K4 using anti-MAP3K4 antibody (M04842). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human U251 whole cell lysates, Lane 2: human K562 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MAP3K4 antigen affinity purified monoclonal antibody (M04842) at 1: 500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for MAP3K4 at approximately 182 kDa. The expected band size for MAP3K4 is at 182 kDa.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-MAP3K4 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.