BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-MAGED1 Antibody (Center)

Catalog Number: M05037

About MAGED1

Involved in the apoptotic response after nerve growth factor (NGF) binding in neuronal cells. Binds NGFR/p75NTR and antagonizes its association with NTRK1/TrkA, inhibits cell cycle progression, and facilitates NGFR-mediated apoptosis. May act as a regulator of the function of DLX family members. May regulate TP53/p53 transcriptional activity and inhibit cell proliferation. Enhances TP53 phosphorylation and accumulation. May enhance ubiquitin ligase activity of RING-type zinc finger-containing E3 ubiquitin-protein ligases. Proposed to act through recruitment and/or stabilization of the Ubl-conjugating enzyme (E2) at the E3:substrate complex.

Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-MAGED1 Antibody (Center)
Reactive Species	Human
Description	Boster Bio Anti-MAGED1 Antibody (Center) (Catalog # M05037). Tested in WB application(s). This antibody reacts with Human.
Application	WB
Clonality	Monoclonal 1305CT862.157.127
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q9Y5V3

Technical Details


Immunogen	This MAGED1 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 20-224 amino acids from the Central region of human MAGED1.
Predicted Reactive Species	Bovine, Rabbit
Isotype	IgG1,kappa

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	WB: 1:1000

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-MAGED1 Antibody (Center) (M05037) Images



Western blot analysis of lysates from Jurkat, 293 cell line (from left to right), using MAGED1 Antibody (Center). M05037 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Anti-MAGED1 Antibody (Center)

For Research Use Only. Not for use in diagnostic procedures.