

## Anti-AMPK beta 2 PRKAB2 Antibody Picoband™ (monoclonal, 6G1)

Catalog Number: M05077

### About PRKAB2

5'-AMP-activated protein kinase subunit beta-2 is an enzyme that in humans is encoded by the PRKAB2 gene. The protein encoded by this gene is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. It is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit may be a positive regulator of AMPK activity. It is highly expressed in skeletal muscle and thus may have tissue-specific roles. Multiple alternatively spliced transcript variants have been found for this gene.

### Overview


Product Name	Anti-AMPK beta 2 PRKAB2 Antibody Picoband™ (monoclonal, 6G1)
Reactive Species	Human
Description	Boster Bio Anti-AMPK beta 2 PRKAB2 Antibody Picoband™ (monoclonal, 6G1) catalog # M05077. Tested in Flow Cytometry, IHC, ICC, WB applications. This antibody reacts with Human.
Application	Flow Cytometry, IHC, ICC, WB
Clonality	Monoclonal 6G1
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	O43741

### Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human AMPK beta 2, different from the related mouse sequence by three amino acids, and from the related rat sequence by two amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit (SV0001-1) for IHC(F) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml</p> <p>Immunohistochemistry (Frozen Section), 0.5-1ug/ml</p> <p>Immunocytochemistry, 0.5-1ug/ml</p> <p>Flow Cytometry, 1-3ug/1x10<sup>6</sup> cells</p>

## Anti-AMPK beta 2 PRKAB2 Antibody Picoband™ (monoclonal, 6G1) (M05077) Images



**Figure 1. Flow Cytometry analysis of PC-3 cells using anti-AMPK beta 2 antibody (M05077).**  
Overlay histogram showing PC-3 cells stained with M05077 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-AMPK beta 2 Antibody (M05077, 1ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-mouse IgG (BA1126, 5-10ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



**Figure 2. Western blot analysis of AMPK beta 2 using anti-AMPK beta 2 antibody (M05077).**  
Electrophoresis was performed on a 10% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysate,  
Lane 2: human placenta tissue lysate,  
Lane 3: human 293T whole cell lysate,  
Lane 4: human A549 whole cell lysate,  
Lane 5: human A375 whole cell lysate,  
Lane 6: human A431 whole cell lysate,  
Lane 7: human U20S whole cell lysate,  
Lane 8: human K562 whole cell lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-AMPK beta 2 antigen affinity purified monoclonal antibody (Catalog # M05077) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.

**Submit a product review to Biocompare.com**

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-AMPK beta 2 PRKAB2 Antibody (monoclonal, 6G1)