

Anti-COX IV COX4I1 Antibody Picoband® (monoclonal, 4G11) Fluoro488 Conjugated

Catalog Number: M05442-1-Fluoro488

About COX4I1

Cytochrome c oxidase subunit 4 isoform 1, mitochondrial is an enzyme that in humans is encoded by the COX4I1 gene. Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer and proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit IV isoform 1 of the human mitochondrial respiratory chain enzyme. It is located at the 3' of the NOC4 (neighbor of COX4) gene in a head-to-head orientation, and shares a promoter with it. Pseudogenes related to this gene are located on chromosomes 13 and 14.

Overview

Product Name	Anti-COX IV COX4I1 Antibody Picoband® (monoclonal, 4G11) Fluoro488 Conjugated
Reactive Species	Human, Mouse, Rat
Application	Flow Cytometry
Clonality	Monoclonal 4G11
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Mouse
Uniprot ID	P13073

Technical Details

Immunogen	E. coli-derived human COX IV recombinant protein (Position: Q59-K169).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Fluoro488

	Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Suggested Dilutions	Flow Cytometry, Optimal dilutions should be determined by end users.

1 Publications Citing This Product

1. PubMed ID: 10.1016/j.bbrc.2017.07.112, Hypoxic postconditioning attenuates apoptosis via inactivation of adenosine A2a receptor through NDRG3-Raf-ERK pathway

Visit bosterbio.com/anti-cox-iv-picoband-trade-antibody-monoclonal-m05442-1-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-COX IV COX4I1 Antibody (monoclonal, 4G11) - Fluoro488

For Research Use Only. Not for use in diagnostic procedures.