

Anti-beta II Tubulin Antibody

Catalog Number: M06868-1

About Tubb2a

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain (By similarity).


Overview

Product Name	Anti-beta II Tubulin Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-beta II Tubulin Antibody (Catalog # M06868-1). Tested in WB, Flow Cytometry, IF application(s). This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IF, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q7TMM9


Technical Details

Immunogen	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 14-46 amino acids from human.
Predicted Reactive Species	Bovine, Chicken, Monkey, Pig, Xenopus
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	IF: 1:25 WB: 1:8000 FC: 1:25


Anti-beta II Tubulin Antibody (M06868-1) Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized C2C12 (mouse myoblast cell line) cells labeling beta II Tubulin with M06868-1 at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on C2C12 cell line. The nuclear counter stain is DAPI (blue).



All lanes : Anti-beta II Tubulin at 1:8000 dilution
Lane 1: mouse brain lysate
Lane 2: Hela whole cell lysate
Lane 3: Jurkat whole cell lysate
Lane 4: PC-12 whole cell lysate
Lane 5: NIH/3T3 whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa
Blocking/Dilution buffer: 5% NFD/MTBST.



Overlay histogram showing NIH/3T3 cells stained with M06868-1 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (M06868-1, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1g/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-beta II Tubulin Antibody

For Research Use Only. Not for use in diagnostic procedures.