

Anti-Hu CD103 Purified ITGAE Monoclonal Antibody

Catalog Number: M07091-1

About ITGAE

CD103 / integrin alphaE is an integrin subunit that is expressed on intraepithelial lymphocytes, epithelial dendritic cells, lamina propria-derived dendritic cells, a subpopulation of lamina propria T cells, a small subset of peripheral lymphocytes, namely T reg cells, and on activated and TGF-beta stimulated lymphocytes. CD103 is in mature form cleaved into two disulfide-linked chains (C-terminal 150 kDa chain and N-terminal 25 kDa chain). It heterodimerizes with integrin beta7 subunit to form alphaE/beta7 integrin (mucosal lymphocyte 1 antigen), which through binding E-cadherin mediates homing of lymphocytes to the intestinal epithelium, and, in addition to the role in adhesion, may serve as an accessory molecule for intraepithelial lymphocyte activation.

Overview

Product Name	Anti-Hu CD103 Purified ITGAE Monoclonal Antibody
Reactive Species	Human, Primate
Description	Boster Bio Anti-Hu CD103 Purified ITGAE Monoclonal Antibody (Catalog# M07091-1). Tested in Flow Cytometry, IP, WB, IHC-F application(s). This antibody reacts with Human, Primate.
Application	Flow Cytometry, IP, IHC-F, WB
Clonality	Monoclonal Ber-ACT8
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage Instructions	Store at 2-8°C. Do not freeze.
Host	Mouse
Uniprot ID	P38570

Technical Details

Immunogen	HTLV-1 induced human T cell line MAPS16. The mouse monoclonal antibody Ber-ACT8 recognizes an extracellular epitope of CD103 (alpha E integrin), a type I transmembrane glycoprotein primarily found on intestinal intraepithelial lymphocytes.
Predicted Reactive Species	Primate
Cross Reactivity	As in case with other anti-CD170 antibodies, this antibody cross-reacts with Siglec-14, whose first two lg domains are almost identical to those of CD170.
Isotype	Mouse IgG1 kappa
Form	Liquid
Concentration	1 mg/ml





Purification	Purified by protein-A affinity chromatography.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Flow cytometry: 1-4 ug/ml.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Hu CD103 Purified ITGAE Monoclonal Antibody