

Anti-Histone H2B (formyl K116) Rabbit Monoclonal Antibody

Catalog Number: M07286-5


Overview

Product Name	Anti-Histone H2B (formyl K116) Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Histone H2B (formyl K116) Rabbit Monoclonal Antibody catalog # M07286-5. Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse.
Application	IF, ICC, WB
Clonality	Monoclonal 31H94
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q16778


Technical Details

Immunogen	A synthesized peptide derived from human Histone H2B (formyl K116)
Isotype	IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 ICC/IF 1:50-200


Anti-Histone H2B (formyl K116) Rabbit Monoclonal Antibody (M07286-5) Images



Western blot analysis of Histone H2B (formyl K116) expression in HeLa cell lysate.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Histone H2B (formyl K116) Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.