

# Anti-AK3L1 (AK4) Mouse Monoclonal Antibody [Clone ID: OTI3B1]

Catalog Number: M07383

#### Overview


| Product Name         | Anti-AK3L1 (AK4) Mouse Monoclonal Antibody [Clone ID: OTI3B1]  |
|----------------------|--|
| Reactive Species     | Dog, Human, Monkey, Mouse, Rat   |
| Description          | Boster Bio AK4 mouse monoclonal antibody, clone OTI3B1 (formerly 3B1). Catalog# M07383. Tested in FC, IF, IHC, WB. This antibody reacts with Human, Monkey, Mouse, Rat, Dog. |
| Application          | Flow Cytometry, IF, IHC, WB  |
| Clonality            | Monoclonal OTI3B1  |
| Formulation          | PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.   |
| Storage Instructions | Store at -20°C as received.  |
| Host                 | Mouse  |
| Uniprot ID           | P27144   |

### **Technical Details**


| Immunogen           | Full length human recombinant protein of human AK4 (NP_037542) produced in HEK293 cell.  |
|---------------------|--|
| Isotype             | lgG1   |
| Concentration       | 1 mg/ml  |
| Purification        | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)  |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows: |




## Anti-AK3L1 (AK4) Mouse Monoclonal Antibody [Clone ID: OTI3B1] (M07383) Images




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY AK4 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-AK4.




Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-AK4 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-AK4 monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer






Carcinoma of Human prostate tissue using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer

Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by







10mM citric buffer




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-AK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer



HEK293T cells transfected with either AK4 (Myc-DDK-tagged) overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-AK4 antibody (M07383)



Anti-AK4 mouse monoclonal antibody (M07383) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY AK4.



Equivalent amounts of cell lysates (10 ug per lane) of wildtype HeLa cells (WT) and AK4-Knockout HeLa cells (KO) were separated by SDS-PAGE and immunoblotted with anti-AK4 monoclonal antibody M07383 (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.

# Submit a product review to Biocompare.com











reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-AK3L1 (AK4) Mouse Monoclonal Antibody [Clone ID: OTI3B1]