

Anti-OVGP1 Antibody (N-Term)

Catalog Number: M07911

About OVGP1

Binds to oocyte zona pellucida in vivo. May play a role in the fertilization process and/or early embryonic development.


Overview

Product Name	Anti-OVGP1 Antibody (N-Term)
Reactive Species	Human
Description	Boster Bio Anti-OVGP1 Antibody (N-Term) (Catalog # M07911). Tested in WB, IHC-P, Flow Cytometry application(s). This antibody reacts with Human.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q12889


Technical Details

Immunogen	This OVGP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 107-137 amino acids from human OVGP1.
Predicted Reactive Species	Human
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:2000 IHC-P: 1:25 FC: 1:25


Anti-OVGP1 Antibody (N-Term) (M07911) Images



All lanes : Anti-OVGP1 Antibody (N-Term) at 1:2000 dilution
Lane 1: human placenta lysate
Lane 2: human uterus lysate
Lane 3: human ovary lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 75 kDa
Blocking/Dilution buffer: 5% NFD/MTBST.



M07911 staining OVGP1 in human fallopian tube tissue sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing Hela cells stained with M07911 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (M07911, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1g/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-OVGP1 Antibody (N-Term)

For Research Use Only. Not for use in diagnostic procedures.