

## Anti-NAT8 Mouse Monoclonal Antibody [Clone ID: OTI5A8]

Catalog Number: M08414


### Overview

Product Name	Anti-NAT8 Mouse Monoclonal Antibody [Clone ID: OTI5A8]
Reactive Species	Human
Description	Boster Bio Anti-NAT8 mouse monoclonal antibody, clone OTI5A8 (formerly 5A8). Catalog# M08414. Tested in IF, IHC, WB. This antibody reacts with Human.
Application	IF, IHC, WB
Clonality	Monoclonal OTI5A8
Formulation	PBS (pH 7.3) containing 1% stabilizing protein, 50% glycerol and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C as received.
Host	Mouse
Uniprot ID	Q9UHE5


### Technical Details

Immunogen	Full-length protein expressed in 293T cell transfected with human NAT8 expression vector
Isotype	IgG1
Concentration	1 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	WB 1:500 IHC 1:50 IF 1:100


## Anti-NAT8 Mouse Monoclonal Antibody [Clone ID: OTI5A8] (M08414) Images




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NAT8 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NAT8.




Immunohistochemical staining of paraffin-embedded Ovary tissue within the normal limits using anti-NAT8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of ovary tissue using anti-NAT8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded pancreas tissue within the normal limits using anti-NAT8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Carcinoma of thyroid tissue using anti-NAT8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded endometrium tissue within the normal limits using anti-NAT8mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Carcinoma of bladder tissue using anti-NAT8mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of breast tissue using anti-NAT8 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded colon tissue within the normal limits using anti-NAT8mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of colon tissue using anti-NAT8mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)

Immunohistochemical staining of paraffin-embedded Kidney tissue within the normal limits using anti-NAT8mouse



monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer)



Anti-NAT8 mouse monoclonal antibody (M08414) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NAT8.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NAT8 Mouse Monoclonal Antibody [Clone ID: OT15A8]

For Research Use Only. Not for use in diagnostic procedures.