

Anti-RGL2 Mouse Monoclonal Antibody [Clone ID: OTI10F2]

Catalog Number: M08483


Overview

Product Name	Anti-RGL2 Mouse Monoclonal Antibody [Clone ID: OTI10F2]
Reactive Species	Human, Mouse, Rat
Description	Boster Bio RGL2 mouse monoclonal antibody, clone OTI10F2. Catalog# M08483. Tested in IHC, WB. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Monoclonal OTI10F2
Formulation	PBS (pH 7.3) containing 1% stabilizing protein, 50% glycerol and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C as received.
Host	Mouse
Uniprot ID	O15211


Technical Details

Immunogen	Human recombinant protein fragment corresponding to amino acids 547-777 of human RGL2 (NP_004752) produced in E.coli.
Isotype	IgG2b
Concentration	1 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	WB 1:2000 IHC 1:150


Anti-RGL2 Mouse Monoclonal Antibody [Clone ID: OTI10F2] (M08483) Images




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RGL2 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RGL2 (1:2000).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human thyroid tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human endometrium tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human prostate tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)


Immunohistochemical staining of paraffin-embedded Human bladder tissue using anti-RGL2 mouse monoclonal antibody.




(Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min



Immunohistochemical staining of paraffin-embedded Human lymph node tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min




Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min




Immunohistochemical staining of paraffin-embedded Human tonsil using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min


Immunohistochemical staining of paraffin-embedded Human colon tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris




buffer (pH8.5) at 120°C for 3min




Immunohistochemical staining of paraffin-embedded Human Kidney tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min



Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min



Immunohistochemical staining of paraffin-embedded Human liver tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min



Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-RGL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min

Submit a product review to [Biocompare.com](https://www.biocompare.com)



Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-RGL2 Mouse Monoclonal Antibody [Clone ID: OTI10F2]

For Research Use Only. Not for use in diagnostic procedures.