

Anti-DPYSL5 Antibody (C-term)

Catalog Number: M08644-1

About DPYSL5

May have a function in neuronal differentiation and/or axon growth.


Overview

| | |
|----------------------|---|
| Product Name | Anti-DPYSL5 Antibody (C-term) |
| Reactive Species | Human, Mouse, Rat |
| Description | Boster Bio Anti-DPYSL5 Antibody (C-term) (Catalog # M08644-1). Tested in WB, IHC-P application(s). This antibody reacts with Human, Mouse, Rat. |
| Application | IHC-P, WB |
| Clonality | Polyclonal |
| Formulation | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. |
| Storage Instructions | Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | Q9BPU6 |


Technical Details

| | |
|----------------------------|---|
| Immunogen | This DPYSL5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 521-554 amino acids from the C-terminal region of human DPYSL5. |
| Predicted Reactive Species | Bovine |
| Isotype | Rabbit IgG |
| Purification | This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Suggested Dilutions | WB: 1:2000 IHC-P: 1:25 |

Anti-DPYSL5 Antibody (C-term) (M08644-1) Images



All lanes : Anti-DPYSL5 Antibody (C-term) at 1:2000 dilution
Lane 1: Y79 whole cell lysates
Lane 2: SH-SY5Y whole cell lysates
Lane 3: Raji whole cell lysates
Lane 4: mouse brain lysates
Lane 5: rat brain lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 61 kDa
Blocking/Dilution buffer: 5% NFD/MTBST.



M08644-1 staining DPYSL5 in human tonsil sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-DPYSL5 Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.