

Anti-UQCRFS1 Mouse Monoclonal Antibody [Clone ID: OTI4H8]

Catalog Number: M08781


Overview

Product Name	Anti-UQCRFS1 Mouse Monoclonal Antibody [Clone ID: OTI4H8]
Reactive Species	Human, Mouse, Rat
Description	Boster Bio UQCRFS1 mouse monoclonal antibody, clone OTI4H8. Catalog# M08781. Tested in IHC, WB. This antibody reacts with Human, Mouse, Rat.
Conjugate	Unconjugated
Application	IHC, WB
Clonality	Monoclonal OTI4H8
Formulation	PBS (pH 7.3) containing 1% stabilizing protein, 50% glycerol and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P47985


Technical Details

Immunogen	Human recombinant protein fragment corresponding to amino acids 79-274 of human UQCRFS1 (NP_005994) produced in E.coli.
Isotype	IgG2a
Concentration	1 mg/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Suggested Dilutions	WB: 1:500~2000 IHC: 1:500


Anti-UQCRFS1 Mouse Monoclonal Antibody [Clone ID: OTI4H8] (M08781) Images




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-UQCRFS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-UQCRFS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-UQCRFS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-UQCRFS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human appendix tissue within the normal limits using anti-UQCRFS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-UQCRFS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)



Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-UQCRFS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Å°C for 3min)



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UQCRFS1 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UQCRFS1 (1:2000).



Western blot analysis of extracts (35ug) from 5 different cell lines and mouse brain tissue lysate by using anti-UQCRFS1 monoclonal antibody (1:500).

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-UQCRFS1 Mouse Monoclonal Antibody [Clone ID: OTI4H8]

For Research Use Only. Not for use in diagnostic procedures.