BOSTER
antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

# Anti-MAGEB4 Mouse Monoclonal Antibody [Clone ID: OTI1B5]

**Catalog Number:** M14698

## Overview

| | |
|---|---|
| Product Name | Anti-MAGEB4 Mouse Monoclonal Antibody [Clone ID: OTI1B5] |
| Reactive Species | Human |
| Description | Boster Bio MAGEB4 mouse monoclonal antibody,clone OTI1B5. Catalog# M14698. Tested in IHC, WB. This antibody reacts with Human. |
| Application | IHC, WB |
| Clonality | Monoclonal OTI1B5 |
| Formulation | PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Storage Instructions | Store at -20°C as received. |
| Host | Mouse |
| Uniprot ID | O15481 |

## Technical Details

| | |
|---|---|
| Immunogen | Human recombinant protein fragment corresponding to amino acids 1-193 of human MAGEB4 (NP_002358) produced in E.coli. |
| Isotype | IgG1 |
| Concentration | 1 mg/ml |
| Purification | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

# Anti-MAGEB4 Mouse Monoclonal Antibody [Clone ID: OTI1B5] (M14698) Images




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAGEB4 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAGEB4 (1:2000).



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-MAGEB4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Â°C for 3min



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-MAGEB4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Â°C for 3min



Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-MAGEB4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120Â°C for 3min