

## Anti-HNRNPA1L2 Antibody (N-Term)

Catalog Number: M16841

### About HNRNPA1L2

Involved in the packaging of pre-mRNA into hnRNP particles, transport of poly(A) mRNA from the nucleus to the cytoplasm and may modulate splice site selection.


### Overview

Product Name	Anti-HNRNPA1L2 Antibody (N-Term)
Reactive Species	Human
Description	Boster Bio Anti-HNRNPA1L2 Antibody (N-Term) (Catalog # M16841). Tested in WB, IHC-P application(s). This antibody reacts with Human.
Application	IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q32P51


### Technical Details

Immunogen	This HNRNPA1L2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 22-56 amino acids from human HNRNPA1L2.
Predicted Reactive Species	Bovine, Chicken, Hamster, Horse, Mouse, Rabbit, Rat
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:2000 IHC-P: 1:25

## Anti-HNRNPA1L2 Antibody (N-Term) (M16841) Images



All lanes : Anti-HNRNPA1L2 Antibody (N-Term) at 1:2000 dilution  
Lane 1: HeLa whole cell lysate  
Lane 2: Jurkat whole cell lysate  
Lane 3: MCF-7 whole cell lysate  
Lane 4: HepG2 whole cell lysate  
Lane 5: K562 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



M16841 staining HNRNPA1L2 in human breast carcinoma tissue sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

### Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HNRNPA1L2 Antibody (N-Term)

For Research Use Only. Not for use in diagnostic procedures.