

## Anti-CDC6 Antibody (Monoclonal, DCS-180)

Catalog Number: MA1019

### About Cdc6

In yeasts, Cdc6 (*Saccharomyces cerevisiae*) and Cdc18 (*Schizosaccharomyces pombe*) associate with the origin recognition complex (ORC) proteins to render cells competent for DNA replication. Cdc6 is overexpressed in human cancers, where it has a critical regulatory role in addition to DNA replication. Yan et al. (1998) showed that Cdc6 is expressed selectively in proliferating but not quiescent mammalian cells, both in culture and within tissues in intact animals. During the transition from a growth-arrested to a proliferative state, transcription of mammalian Cdc6 is regulated by E2F proteins, as revealed by a functional analysis of the human Cdc6 promoter and by the ability of exogenously expressed E2F proteins to stimulate the endogenous Cdc6 gene. They conclude that expression of human Cdc6 is regulated in response to mitogenic signals through transcriptional control mechanisms involving E2F proteins, and that Cdc6 is required for initiation of DNA replication in mammalian cells.

### Overview

Product Name	Anti-CDC6 Antibody (Monoclonal, DCS-180)
Reactive Species	Human
Description	Boster Bio Anti-CDC6 Antibody (Monoclonal, DCS-180) catalog # MA1019. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Monoclonal DCS-180
Formulation	Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN <sub>3</sub> as preservative.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	O89033

### Technical Details

Immunogen	Recombinant human Cdc6.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Mouse IgG1
Form	Lyophilized
Concentration	Adding 1 ml of PBS buffer will yield a concentration of 100 ug/ml.

Purification	Ascites
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used: Western blot, 1-2ug/ml, Human</p>

## Anti-CDC6 Antibody (Monoclonal, DCS-180) (MA1019) Images

---



Boster Kit Box

### Submit a product review to Biocompare.com

---

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CDC6 Antibody (Monoclonal, DCS-180)