

Anti-p19INK4d Cdkn2d Antibody (Monoclonal, DCS-100)

Catalog Number: MA1075

About Cdkn2d

Cyclins are important in regulating the cell cycle through their formation of enzymatic complexes with various cyclin-dependent kinases. P19 (INK4d) also known as cyclin-dependent kinase inhibitor 2D, is one of the novel members of the mouse INK4 gene family. Okuda et al. (1995) described the cloning of the human INK4d gene (CDKN2D). The predicted 166-amino acid protein is 86% identical to the mouse protein and about 45% identical to other human INK4 family members.

Overview


Product Name	Anti-p19INK4d Cdkn2d Antibody (Monoclonal, DCS-100)
Reactive Species	Human
Description	Boster Bio Anti-p19INK4d Cdkn2d Antibody (Monoclonal, DCS-100) catalog # MA1075. Tested in IHC, ICC, WB applications. This antibody reacts with Human.
Application	IHC, ICC, WB
Clonality	Monoclonal DCS-100
Formulation	Mouse IgG in stabilizing components, 1.2% sodium acetate and 0.01mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q60773

Technical Details

Immunogen	Recombinant human p19INK4d.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit (SV0001-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Mouse IgG1
Form	Lyophilized
Concentration	Adding 1 ml of PBS buffer will yield a concentration of 100 ug/ml.
Purification	Ascites
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 1-2ug/ml, Human

Immunocytochemistry , 1ug/ml, Human, -
Western blot, 0.5-1ug/ml, Human

Anti-p19INK4d Cdkn2d Antibody (Monoclonal, DCS-100) (MA1075) Images



Western blot analysis of p19INK4d using anti-p19INK4d antibody (MA1075). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human MOLT-4 whole cell lysates, Lane 2: human HEL whole cell lysates, Lane 3: human Jurkat whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-p19INK4d antigen affinity purified monoclonal antibody (Catalog # MA1075) at 1 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for p19INK4d at approximately 18 kDa. The expected band size for p19INK4d is at 18 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-p19INK4d Cdkn2d Antibody (Monoclonal, DCS-100)

For Research Use Only. Not for use in diagnostic procedures.