

Anti-Phospho-Tau (T231) MAPT Rabbit Monoclonal Antibody

Catalog Number: P00097-1

Overview

Product Name	Anti-Phospho-Tau (T231) MAPT Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Phospho-Tau (T231) MAPT Rabbit Monoclonal Antibody catalog # P00097-1. Tested in WB, IHC, IP applications. This antibody reacts with Human, Mouse, Rat.
Application	IP, IHC, WB
Clonality	Monoclonal EIH-13
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P10636

Technical Details

Immunogen	A synthesized peptide derived from human Phospho-Tau (T231)
Predicted Reactive Species	Human, Primate
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>WB 1:500-1:2000</p> <p>IHC 1:50-1:200</p> <p>IP 1:50</p>

Anti-Phospho-Tau (T231) MAPT Rabbit Monoclonal Antibody (P00097-1) Images




Figure 1. Western blot analysis of Tau using anti-Tau antibody (P00097-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat brain tissue lysates,

Lane 2: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Tau antigen affinity purified monoclonal antibody (Catalog # P00097-1) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Tau at approximately 50-70 kDa. The expected band size for Tau is at 79 kDa.




Figure 2. IHC analysis of Tau using anti-Tau antibody (P00097-1).

Tau was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Tau Antibody (P00097-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.





Figure 3. IHC analysis of Tau using anti-Tau antibody (P00097-1).

Tau was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Tau Antibody (P00097-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of Phospho-Tau (T231) expression in (1) SH-SY5Y cell lysate; (2) SH-SY5Y cell lysate, treated with sorbitol.

6 Publications Citing This Product

1. PubMed ID: 28934273, Han F1, Zhuang TT1, Chen JJ1, Zhu XL1,2, Cai YF1, Lu YP1. PLoS One. 2017 Sep 21;12(9):e0185102. doi: 10.1371/journal.pone.0185102. eCollection 2017. Novel derivative of Paeonol, Paeononlsilatie sodium, alleviates behavioral damage and hippocampal ...
2. PubMed ID: 28123476, The expression of insulin-like growth factor-1 in senior patients with diabetes and dementia
3. PubMed ID: 30102687, Micro-RNA-137 Inhibits Tau Hyperphosphorylation in Alzheimer's Disease and Targets the CACNA1C Gene in Transgenic Mice and Human Neuroblastoma %u2026

Visit bosterbio.com/anti-phospho-tau-t231-rabbit-monoclonal-antibody-p00097-1-boster.html to see all 6 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Phospho-Tau (T231) MAPT Rabbit Monoclonal Antibody