

Anti-Phospho-NFATC2 (S326) Antibody

Catalog Number: P00969

About NFATC2

NFATC2(Nuclear factor of activated T-cells, cytoplasmic 2), also known as NFATP or the 'preexisting component' of NFAT, is present in the cytosolic fraction of unstimulated T cells, which is also a member of the nuclear factor of activated T cells (NFAT) family. The NFATC2 gene is mapped on 20q13.2. NFATC2 is highly homologous to NFATC1 over a limited domain which shows similarity to the Dorsal/Rel family but has a wider tissue distribution. Ectopic expression of NFATC2 inhibited the basal activity of the human CDK4 promoter. Additionally, both Calna^{-/-} and Nfatc2^{-/-} mice had elevated protein levels of Cdk4, confirming a negative regulatory role for the calcineurin/NFAT pathway. NFATC2 controls myoblast fusion at a specific stage of myogenesis after the initial formation of a myotube and is necessary for further cell growth. Overexpression of NFATC2 promoted differentiation of osteoclast precursor cells into tartrate-resistant acid phosphatase-positive (TRAP-positive) multinucleated osteoclast-like cells even in the absence of RANKL.

Overview

Product Name	Anti-Phospho-NFATC2 (S326) Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Phospho-NFATC2 (S326) Antibody catalog # P00969. Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg stabilizing protein and 50% glycerol This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	12 months from date of receipt at -20°C as supplied. 6 months at 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q13469

Technical Details

Immunogen	A synthesized peptide derived from human NFAT1 around the phosphorylation site of Ser326.
Form	Liquid
Concentration	500 ug/ml
Purification	Protein A affinity purified.

Suggested Dilutions

Western blot, 1:500-2000
Immunohistochemistry, 1:50-200

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Phospho-NFATC2 (S326) Antibody

For Research Use Only. Not for use in diagnostic procedures.