

Anti-CXCR2 Antibody

Catalog Number: PA1029

About CXCR2

CXCR2 is a receptor for Interleukin 8, which is a powerful neutrophil chemotactic factor. It is a member of the GPCR family (subfamily, chemokine). Binding of IL8 to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activate a phosphatidylinositol-calcium second messenger system. This receptor binds to IL8 with a high affinity and to GRO/MGSA and NAP2 also with a high affinity. It has been reported to be expressed in a wide variety of tissues. ESTs have been isolated from human placenta and thymus libraries.

Overview

Product Name	Anti-CXCR2 Antibody
Reactive Species	Human, Rat
Description	Boster Bio Anti-CXCR2 Antibody catalog # PA1029. Tested in WB applications. This antibody reacts with Human, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P25025

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human CXCR2, different from the related mouse sequence by six amino acids.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human, Rat</p>

Anti-CXCR2 Antibody (PA1029) Images




Figure 1. Western blot analysis of CXCR2 using anti-CXCR2 antibody (PA1029).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat lung tissue lysates,

Lane 2: human COLO320 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CXCR2 antigen affinity purified polyclonal antibody (Catalog # PA1029) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CXCR2 at approximately 45 kDa. The expected band size for CXCR2 is at 41 kDa.

7 Publications Citing This Product

1. PubMed ID: 30404567, Cui D,Zhao Y,Xu J.Activation of CXCL5-CXCR2 axis promotes proliferation and accelerates G1 to S phase transition of papillary thyroid carcinoma cells and activates JNK and p38 pathways.Cancer Biol

Ther.2019;20(5):608-616.doi:10.1080/15384047.2018.1539289.Epub 2018 Nov 7.PMID:30404567;PMCID:PMC6606038.

2. PubMed ID: 24852102, Chemokine CXCL1 enhances inflammatory pain and increases NMDA receptor activity and COX-2 expression in spinal cord neurons via activation of CXCR2

3. PubMed ID: 23831863, Zhang Zj, Cao DI, Zhang X, Ji Rr, Gao Yj. Pain. 2013 Oct;154(10):2185-97. Doi: 10.1016/J.Pain.2013.07.002. Epub 2013 Jul 4. Chemokine Contribution To Neuropathic Pain: Respective Induction Of Cxcl1 And Cxcr2 In Spinal Cord Astrocytes And Neurons.

Visit bosterbio.com/anti-cxcr2-antibody-pa1029-boster.html to see all 7 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CXCR2 Antibody