

Anti-VEGF/VEGFA Antibody Picoband®

Catalog Number: PA1080

About VEGFA

VEGF, a homodimeric glycoprotein of relative molecular mass 45,000, is the only mitogen that specifically acts on endothelial cells. It may be a major regulator of tumor angiogenesis in vivo. Vascular endothelial growth factor is a mitogen primarily for vascular endothelial cells. It is, however, structurally related to platelet-derived growth factor. VEGF shares homology with the PDGF A chain and B chain, including conservation of all 8 cysteines found in PDGFA and PDGFB. VEGF gene contains 8 exons. Vascular endothelial growth factor (VEGF) induces remodeling and enhances TH2-mediated sensitization and inflammation in the lung. VEGF regulates haematopoietic stem cell survival by an internal autocrine loop mechanism. Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo.

Overview


Product Name	Anti-VEGF/VEGFA Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-VEGF/VEGFA Antibody catalog # PA1080. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.01mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P15692

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human VEGF, identical to the related mouse and rat sequences.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western

	blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human


Anti-VEGF/VEGFA Antibody Picoband® (PA1080) Images



Western blot analysis of VEGFA using anti-VEGFA antibody (PA1080). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HeLa whole cell lysates, Lane 2: human MCF-7 whole cell lysates, Lane 3: rat kidney tissue lysates, Lane 4: mouse kidney tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-VEGFA antigen affinity purified polyclonal antibody (Catalog # PA1080) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for VEGFA at approximately 27 kDa. The expected band size for VEGFA is at 27 kDa.




IHC analysis of VEGFA using anti-VEGFA antibody (PA1080). VEGFA was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-VEGFA Antibody (PA1080) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of VEGFA using anti-VEGFA antibody (PA1080). VEGFA was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-VEGFA Antibody (PA1080) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.


Molecular mechanism validation of P-Nb2C-PEG@si-circPUM1 anti-angiogenic performance in vivo. (A) H&E staining showed increased necrotic loci within tumors of P-Nb2C-PEG@si-circPUM1 group. (B, D) IHC staining of Ki-67




demonstrated a clear reduction in Ki-67-positive cells in P-Nb2C-PEG@si-circPUM1 group. (C, F) IHC staining of CD31 indicated a marked decrease in MVD within tumors of P-Nb2C-PEG@si-circPUM1 group. (H) QPCR showed a significant downregulation of circPUM1 expression in the P-Nb2C-PEG@si-circPUM1 treated group. IHC (E, G) and Western blot (I, J) demonstrated significant down-regulation of RAB27B and VEGFA. Index in PubMed under a CC BY license. PMID: 41050095



ZGF Promotes VEGF/Notch1/Noggin Pathway Activation. (A) RT-PCR was used to detect NOG, Notch1, and VEGF mRNA expression levels in the tibial metaphysis of Wistar and GK rats. (B-C) Target proteins NOG, Notch1, and VEGF expression levels were measured by Western blot in the tibial metaphysis of Wistar and GK rats. (Uncropped Western blot images are included in Additional file 1.) Data are expressed as mean \pm SD. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001 VS. MC group. #P < 0.05, ##P < 0.01, ###P < 0.001, ####P < 0.0001 VS. PC group. ^P < 0.05, ^^P < 0.01, ^^P < 0.001, ^^P < 0.0001 VS. BC group, (n = 3). Index in PubMed under a CC BY license. PMID: 38524608




In vitro antitumor performance of P-Nb2C@si-circPUM1 in ovarian cancer cells. (A) Confocal microscopy images showed intracellular uptake of Cy3-labeled circPUM1 siRNA. Cell apoptosis assays (B), Transwell assays (C), wound healing assays (E) and EdU proliferation assays (F) showed that P-Nb2C-loaded circPUM1 siRNA effectively suppressed ovarian cancer cell phenotypes. (G) RT-qPCR demonstrated a significant downregulation of circPUM1 expression in lipo2000-transfected and P-Nb2C-loaded siRNA groups. Immunofluorescence (D) and Western blot (H) validated the downregulation of VEGFA and RAB27B expression in cells treated with P-Nb2C@si-circPUM1. (I) CCK8 assays showed inhibition of ovarian cancer cell viability in lipo2000-transfected and P-Nb2C-loaded siRNA groups. Index in PubMed under a CC BY license. PMID: 41050095




The anti-angiogenic activity of SPS on HUVECs in vitro. (a) Dose- and time-dependent growth-inhibiting effects of SPS on HUVECs. Cells were cultured in 96-well plate and treated with different concentrations of SPS (0-1.5 mg/ml) for 12 and 24 h. The cell viability was analyzed by MTT assay. Each data indicated the mean \pm SD of three independent experiments (n = 3). * P


Western blot analysis of the human and chicken VEGF-A protein in the CAM. In the NCI-H446/HIF-1alpha and NCI-H446/siHIF-1alpha groups, the SCLC cells were transduced




with Ad-HIF-1alpha or Ad-siHIF-1alpha (MOI = 50) for 60 h before implanting onto the CAM to form transplantation tumors. Western blots were performed to detect the VEGF-A protein level in the tumors and peripheral tissues on day 17 of incubation. Data are presented as means \pm SD. (A) Representative images of three independent experiments (Lane A - human VEGF-A protein expression in the tumors from the NCI-H446 group; Lane B - human VEGF-A protein expression in the tumors from the NCI-H446/HIF-1alpha group; and Lane C - human VEGF-A protein expression in the tumors from the NCI-H446/siHIF-1alpha group) (human - * $p < 0.05$ group C vs. group B; ** $p < 0.05$ group C vs. group D) (chicken - * $p < 0.05$ group C vs. group B; ** $p < 0.05$ group C vs. group D). (B) Representative images of three independent experiments (Lane A - chicken VEGF-A protein expression of control group; Lane B - chicken VEGF-A protein expression in the tumors from the NCI-H446 group; Lane C - chicken VEGF-A protein expression in the tumors from the NCI-H446/HIF-1alpha group; and Lane D - Chicken VEGF-A protein expression in tumors from the NCI-H446/siHIF-1alpha group). (C) Densitometry analysis of the relative expression of VEGF-A protein compared to the corresponding beta-actin in each group ($p < 0.05$). Index in PubMed under a CC BY license. PMID: 21843314




STAT1 inhibitor attenuated IL-17-induced IL-6, IL-8, and VEGF production in human lung adenocarcinoma. A549/H292 cells were incubated with IL-17 or IL-17 plus a STAT1 inhibitor for 6 or 48 h (100 ng/ml IL-17; and 30 μ M inhibitor). The IL-6, IL-8, and VEGF mRNA and protein levels were determined by qRT-PCR and ELISA, respectively. (A) IL-6, IL-8, and VEGF mRNA levels in A549/H292 cells, mRNA expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method, and target gene expression was normalized to the GAPDH housekeeping gene. The data are presented as the mean \pm SEM of three independent experiments. (B) IL-6, IL-8, and VEGF protein levels in A549/H292 cells. The data are presented as the mean \pm SEM of three independent experiments. Comparisons were performed using the t-test. * $p < 0.05$; ** $p < 0.01$; and *** $p < 0.001$. Index in PubMed under a CC BY license. PMID: 27819281




IL-6, IL-8, VEGF and STAT1 expression were augmented in A549-IL-17 cell-bearing nude tissues. (A, B) The relative protein expression of IL-6, IL-8, VEGF and STAT1 in tumour tissues of A549-IL-17 vs. A549-Neo cell-bearing nude mice was determined by WB. (C) The relative mRNA expression of IL-6, IL-8, VEGF and STAT1 in tumour tissues of A549-IL-17 vs. A549-Neo cell-bearing nude mice was determined by qRT-PCR ($n = 5$). mRNA expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method, and target gene expression was normalized to the GAPDH housekeeping gene. The data are presented as the mean \pm SEM for five mice per group, and the results are representative of two independent experiments. * $p < 0.05$; ** $p < 0.01$; and *** $p < 0.001$. Index in PubMed under a CC BY license. PMID: 27819281




Expression of CD31, IL-17, IL-6, IL-8, and VEGF protein in human lung adenocarcinoma tissues. Immunohistochemical determination of CD31, IL-17, IL-6, IL-8, VEGF protein in 30 patients with lung adenocarcinoma. (A - F) High CD31, IL-17, IL-6, IL-8, VEGF protein expression were presented in tumor tissue of case 58. (G - L) Low CD31, IL-17, IL-6, IL-8, VEGF protein expression were showed in tumor tissue of case 34. No staining was observed when an isotype-matched control mAb (A, G) was used (magnification, $\times 200$). Index in PubMed under a CC BY license. PMID: 27819281



Correlation between MVD and IL-17, IL-6, IL-8, VEGF in human lung adenocarcinoma tissues. IL-17, IL-6, IL-8, VEGF mRNA and protein and CD31 protein levels were determined in human lung adenocarcinoma tissues by qRT-PCR or IHC, respectively. (A) Spearman's correlation analysis was performed to analyse the correlation between IL-17 (a), IL-6 (b), IL-8 (c), VEGF (d) protein expression and tumour microvessel density (MVD) by CD31 staining in tumor tissues with human lung adenocarcinoma. (B) Spearman's correlation analysis was performed to analyse the correlation between IL-17 (a), IL-6 (b), IL-8 (c), VEGF (d) expression and tumour microvessel density by CD31 staining in 28 tissues with human lung adenocarcinoma. mRNA expression levels were calculated using the $-\Delta\Delta Ct$ method, and target gene expression was normalized to the GAPDH housekeeping gene. Index in PubMed under a CC BY license. PMID: 27819281




Vascular endothelial growth factor (VEGF) ligands and receptors were found to be essential for the angiogenesis-promoting role of deer TMSB10 in HUVECs. a Representative images of VEGFA, VEGFB, VEGFC, VEGFD, FLT1, KDR, and FLT4 protein levels in HUVECs overexpressing deer TMSB10 after Western blot analysis. b Quantification of Western blotting for the VEGF family. The protein levels were normalized with GAPDH. The relative levels of proteins were calculated and plotted. The data are expressed as mean \pm SD, n = 3 experiments. * P




Effects of deer TMSB10 on cell proliferation, tube formation, and motility of HUVECs as well as on DRG outgrowth. a HUVECs overexpressing deer TMSB10 compared with vector alone. HUVECs overexpressing deer TMSB10 had significantly higher growth rates than that of the vector control group. b Determination of HUVEC proliferation using the MTT assay. HUVEC cells were exposed for 24, 48, and 72 h in the presence of deer exogenous TMSB10 (eTMSB10: 50, 100, or 150 ng/ml) and compared with control. The growth rates of the HUVECs treated with deer eTMSB10 (50, 100, and 150 ng/ml) were significantly higher than in controls. c Representative images (inverted phase contrast) of tube formation assays. Scale bars = 200 μ m. d Quantification of tube formation by calculating the average number of branched vessels per field of view in response to exogenous TMSB10 (eTMSB10: 50, 100, or 150 ng/ml),


overexpressed TMSB10 (TMSB10), and vascular endothelial growth factor (VEGF) as a positive control. Results showed that deer eTMSB10 and deer TMSB10 overexpressing HUVECs showed more tube formation in the HUVECs compared with controls or vector. e Transwell migration assays with HUVECs treated with deer eTMSB10 (50, 100, and 150 ng/ml) or overexpressing deer TMSB10 were compared with their controls. Results showed that deer eTMSB10 and deer TMSB10 overexpressing HUVECs showed more migration compared with the control or vector. f Lamellipodium emerging from a DRG neuron after treatment for 2 days or 5 days with 50, 100, and 150 ng/ml eTMSB10 or nerve growth factor (NGF; 50 ng/ml) as a positive control. Scale bar = 100 μ m. g Quantification of average length of lamellipodium. Results showed that deer eTMSB10 significantly increased the growth of lamellipodium from DRG compared with the control. Data represent mean \pm SD of three or four experiments. * P




Effect of miR-542-3p overexpression on angiogenic capacity of HESCs. (a) q-PCR analysis of VEGF, MMP9, COX2 mRNA levels in control or miR-542-3p mimic or mimics NC groups. (b) Representative western blots showing VEGF, MMP9, COX2 protein levels in control or miR-542-3p mimic or mimics NC groups. (c) Capillary-like tube formation performed by HUVECs with /without the supernatant of miR-542-3p-overexpressing HESCs. Representative images (100X) are shown. Number of Tube area (tube meshes and total master segment length) were acquired automatically. * $p < 0.05$, ** $p < 0.01$. Values are expressed as mean \pm SD of three independent experiments. Index in PubMed under a CC BY license. PMID: 33785768



Transplantation of PDGF-BB-primed MSCs via UTM reduces cardiomyocyte apoptosis and improves angiogenesis in rat hearts post-MI. (A) Representative images of TUNEL-positive cardiomyocytes in the ischemic area 30 days after MI. Apoptotic nuclei were identified as TUNEL positive (green fluorescence), and total nuclei were identified by DAPI counterstaining (blue fluorescence). Myocardium was stained using a monoclonal antibody against cardiac troponin I (red fluorescent). Bar, 20 μ m. (B) Quantification of TUNEL-positive cardiomyocytes. (C - F) Western blotting of activated caspase 3, Bax, and BCL-2 in the ischemic heart. GAPDH was used as a loading control. (G) Representative images of CD31 staining and alpha-SMA staining in the ischemic hearts of rats 30 days post-MI. Bar, 20 μ m. (H) Quantitative analysis of the capillary density in the ischemic heart. (I) Quantitative analysis of the arteriole density in the ischemic heart. (J - M) Protein expression of VEGF, bFGF and IGF-1 determined by Western blotting in ischemic myocardium, with GAPDH as the internal control. Values are the mean \pm SEM. Significant differences were determined by using one-way ANOVA. N = 5/group. * $p < 0.01$ vs. MI; # $p < 0.01$ vs. MI-MSC-vehicle Index in PubMed under a CC BY license. PMID: 38102643




As 2 O 3 inhibited angiogenesis by modulating TSP-1, TGF-beta1, CTGF and VEGF expression in RA-FLS. RA-FLS and HDMECs co-cultures were respectively treated with As 2 O 3 alone or together with TNF-alpha (100ng/ml) for 48 h. A. TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and HDMECs co-cultures were analyzed by ELISA. Results showed that concentrations of TSP-1, TGF-beta1, CTGF and VEGF increased significantly after treatment with TNF-alpha (100 ng/ml) (n = 3) compared to vehicle control group (n = 3; & p < 0.05), and then significantly decreased concentrations of TSP-1, TGF-beta1, CTGF and VEGF were observed after the treatment of As 2 O 3 at doses of 1.0 uM and 2.0 uM (n = 3, respectively; # p < 0.05, ## p < 0.01). TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and HDMECs co-cultures without TNF-alpha addition also decreased significantly after treatment of As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) compared to vehicle control (n = 3; * p < 0.05). B. mRNA levels of TSP1, TGF-beta1, CTGF and VEGF expression in RA-FLS co-cultured were performed by real-time PCR. Results showed similar changes of TSP-1, TGF-beta1, CTGF and VEGF mRNA expression as demonstrated in protein regulation after treatment of As 2 O 3 alone (n = 3) or together with TNF-alpha (100ng/ml) (n = 3; * p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01). C. and D. Transwell assay and tube formation test were performed by applying supernatants from RA-FLS and HDMECs co-culture to HDMECs for 6 h respectively. Results showed that migration and capillary-like structure formation of HDMECs significantly increased after TNF-alpha (100ng/ml) stimulation (n = 3, respectively; * p < 0.05), while As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) played a significantly opposite role in migration and tube formation with or without TNF-alpha (* p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01), which were also in a dose dependent manner. E. Ex vivo aortic ring angiogenesis assay showed similar changes of microvessel sprouting as demonstrated in migration and tube formation of HDMECs in the transwell assay and tube formation test above (n = 3, respectively; * p < 0.05, ** p < 0.01, # p < 0.05). Bars = 300um. Original magnification = x5. Results are expressed as the mean ± S.E.M. & p < 0.05 versus Veh. * p < 0.05, ** p < 0.01 versus Veh, # p < 0.05, ## p < 0.01 versus TNF-alpha. Veh = vehicle control under treatment of 1% FBS DMEM alone. TNF-alpha = control group under treatment of TNF-alpha (100ng/ml).Index in PubMed under a CC BY license. PMID: 29088724




Increased expression of TSP-1, TGF-beta1, CTGF and VEGF in supernatants of RA-FLS and HDMECs co-culture compared to NH-FLS and HDMECs co-culture. Normal human (NH) FLS and rheumatoid arthritis (RA) FLS were co-cultured with HDMECs for 48 h, respectively. A. ELISA analysis demonstrated significant increase in the concentrations of TSP-1, TGF-beta1, CTGF and VEGF in supernatants of RA-FLS and HDMECs co-culture (n = 3) compared with those from NH-FLS and HDMECs co-culture (n = 3; p < 0.05). B. Real-time PCR analysis showed increased mRNA expression of TSP-1, TGF-beta1, CTGF and VEGF in RA-FLS co-cultured (n = 3) compared to NH-FLS co-cultured (n = 3; * p < 0.05, ** p <

0.01, *** $p < 0.001$). C. and D. Transwell assay (C; $n = 3$) and tube formation test (D; $n = 3$) for 6 h demonstrated significant up-regulation in migration and capillary-like structure formation of HDMECs respectively under treatment of supernatants from RA-FLS and HDMECs co-culture ($n = 3$) compared to those from NH-FLS and HDMECs co-culture ($n = 3$; * $p < 0.05$). E. aortic rings were placed on GFR-Matrigel-coated plates and incubated in 1% FBS EGM-2. On 3rd day, the EGM-2 were exchanged with supernatants from FLS and HDMECs co-culture and further incubated for 3 days. Ex vivo aortic ring angiogenesis assay showed significant up-regulation in microvessel sprouting under treatment of supernatants from RA-FLS and HDMECs co-culture ($n = 3$) compared to those from NH-FLS and HDMECs co-culture ($n = 3$; * $p < 0.05$). Bars = 300 μ m. Original magnification = $\times 5$. Results are expressed as the mean \pm S.E.M. Index in PubMed under a CC BY license. PMID: 29088724




Schematic representation of PDGF-BB-primed MSC transplantation by UTMD for infarcted myocardium repair. UTMD enhanced the delivery of MSCs into the infarcted myocardium by upregulating SDF-1 expression. Compared with UTMD alone, UTMD combined with PDGF-BB pretreatment increased the therapeutic effect of grafted cells by improving MSC migration and survival, reducing cardiomyocyte apoptosis, decreasing fibrosis, increasing microvessel density (via upregulation of VEGF, bFGF and IGF-1) and improving cardiac function. PDGF-BB promotes the survival/retention and cardioprotection of engrafted MSCs in rat models of MI via the PI3K/Akt pathway and CXCR4 activation. Index in PubMed under a CC BY license. PMID: 38102643




Correlation between IL-17 and IL-6, IL-8, VEGF in human lung adenocarcinoma tissues. IL-17, IL-6, IL-8, VEGF mRNA and protein levels were determined in human lung adenocarcinoma tissues by qRT-PCR or IHC, respectively. (A) Spearman's correlation analysis was performed to analyse the correlation between IL-17 protein expression and IL-6 (a), IL-8 (b), VEGF (c) protein in tumour tissues of patients with lung adenocarcinoma. (B) Pearson's correlation analysis was used to analyse the relationship between IL-17 mRNA expression and IL-6 (a), IL-8 (b), VEGF (c) mRNA expression in tumour tissues of patients with lung adenocarcinoma. mRNA expression levels were calculated using the $-\Delta\Delta Ct$ method, and target gene expression was normalized to the GAPDH housekeeping gene. Index in PubMed under a CC BY license. PMID: 27819281


STAT1 siRNA reversed IL-17-induced IL-6, IL-8, and VEGF expression in human lung adenocarcinoma cells in vitro. A549/H292 cells (2×10^5 cells/well) were transfected with STAT1 siRNA for 48 h. (A) STAT1 protein expression was determined by WB. Then, A549/H292 cells (5×10^5 cells/well) were transfected with siRNA for 48 h and subsequently treated with human IL-17 (100 ng/ml) for an additional 6 or 48 h. The IL-6, IL-8, and VEGF mRNA and



protein expression levels were determined by qRT-PCR and WB, respectively. (B) IL-6, IL-8, VEGF mRNA levels in A549/H292 cells. mRNA expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method, and target gene expression was normalized to the GAPDH housekeeping gene. The data are presented as the mean \pm SEM of three independent experiments. (C) IL-6, IL-8, and VEGF protein levels in A549/H292 cells and the results are presented as the mean \pm SEM of three independent experiments. Comparisons were performed using the t-test. * $p < 0.05$; ** $p < 0.01$; and *** $p < 0.001$. Index in PubMed under a CC BY license. PMID: 27819281




The modulation of TSP-1, TGF-beta1, CTGF and VEGF expression alongside interventions of TSP-1, TGF-beta1, CTGF and VEGF stimulation or knockdown. A. The effects of TSP-1, TGF-beta1, CTGF and VEGF stimulation/silencing in RA-FLS on TSP-1, TGF-beta1, CTGF, and VEGF proteins expression in the supernatants were determined by ELISA. RA-FLS were stimulated with CTGF (500 ng/ml), TGF-beta1 (5ng/ml), TSP-1 (1,000 ng/ml) and VEGF (50 ng/ml) respectively for 48 h. Stimulation (n = 3)/silencing (n = 3) of TGF-beta1 in RA-FLS resulted in significant increase/decrease in the supernatants protein expression of TSP-1, CTGF, and VEGF (p < 0.05). Stimulation (n = 3)/silencing (n = 3) of VEGF in RA-FLS led to similar changes of the other three protein expression in supernatants (p < 0.05). Addition(n = 3)/silencing(n = 3) of CTGF in RA-FLS caused significant up-/down-regulation in the supernatants protein expression of TSP-1 and VEGF (p < 0.05). And stimulation (n = 3)/knockdown (n = 3) of TSP-1 in RA-FLS didn't lead to significant increase/decrease in the expression of supernatants protein TGF-beta1, CTGF and VEGF. B. TSP-1, TGF-beta1, CTGF and VEGF mRNA expression levels were analyzed by real-time PCR after TSP-1, TGF-beta1, CTGF and VEGF stimulation/silencing. Results showed similar alterations of TSP-1, TGF-beta1, CTGF and VEGF mRNA expression as demonstrated in supernatants protein regulation after TSP-1, TGF-beta1, CTGF and VEGF stimulation/silencing (n = 3 respectively; p < 0.05). Results are expressed as the mean \pm S.E.M. * p < 0.05, ** p < 0.01, *** p < 0.001 versus Veh, # p < 0.05, ## p < 0.01, ### p < 0.001 versus Scr siRNA. Veh = vehicle control. Scr siRNA = scramble siRNA. Index in PubMed under a CC BY license. PMID: 29088724




Increased VEGF and VEGFR2 expression in tumors from CD47-deficient mice. (A) Representative images of tumor sections stained for VEGF-A (brown). Scale bar, 20 μ m. (B) Percentages of VEGFR2+ area by counting positive cells in six randomly selected fields (400 \times) using Image Pro Plus 6.0 software. (C) mRNA levels of VEGF-A in tumor samples from WT and CD47^{-/-} mice (n = 4 per group). (D) Representative images of tumor sections stained for VEGFR2+ (brown). Scale bar, 20 μ m. (E) Percentages of VEGFR2+ area by counting positive cells in six randomly selected fields (400 \times) using Image Pro Plus 6.0 software. (F) mRNA levels of VEGFR2 in tumor samples from WT and CD47^{-/-}


mice (n = 4 per group). Data are mean ± SDs. * P < 0.05, ** P < 0.01; *** P < 0.001. Index in PubMed under a CC BY license. PMID: 27283989



The effect of IL-17 on IL-6, IL-8, VEGF expression in human lung adenocarcinoma cells in vitro . A549/H292 cells were incubated with IL-17 or IL-17 (100 ng/ml) for 6 or 48 h. The IL-6, IL-8, and VEGF mRNA and protein levels were determined by qRT-PCR or ELISA, respectively. (A) IL-6, IL-8, and VEGF mRNA levels in A549/H292 cells mRNA expression levels were calculated using the 2^{-ΔΔCt} method, and target gene expression was normalized to the GAPDH housekeeping gene. The data are presented as the mean ± SEM of three independent experiments. (B) IL-6, IL-8, and VEGF protein levels in A549/H292 cells by ELISA. The results shown are representative of four independent experiments and are presented the mean ± SEM. Comparisons were performed using the t-test. *p<0.05; **p<0.01; and ***p<0.001. Index in PubMed under a CC BY license. PMID: 27819281




RT-PCR analysis of human and chicken angiogenic factors mRNA . Microarray analysis was performed to screen out the angiogenic factors affected by HIF-1alpha in SCLC cells (table 2). Afterwards, RT-PCR analysis was used to detect the expression of angiogenic factors affected by HIF-1a in the transplantation tumors of CAM in vivo . (A), Human and chicken VEGF-A, TNFAIP6, PDGFC, FN1, MMP28, MMP14, SOCS2 and IGFBP3 mRNA expression: Representative images of three independent experiments (Lane 1: control group-no human mRNA expression, Lane 2: transplantation tumor of NCI-H446 cells transduction by empty vector Ad5-NCI-H446 cells group, Lane 3: ransplantation tumor of NCI-H446 cells with transduction by HIF-1alpha-NCI-H446/HIF-1alpha group, Lane 4: transplantation tumor of NCI-H446 cells with transduction by siHIF-1alpha-NCI-H446/siHIF-1alpha group). (B and C), Relative expression levels of mRNA in NCI-H446/HIF-1alpha group and NCI-H446/siHIF-1alpha group compared with that in control group and NCI-H446 cells group (p < 0.05). Index in PubMed under a CC BY license. PMID: 21843314




As 2 O 3 inhibited angiogenesis by modulating TSP-1, TGF-beta1, CTGF and VEGF expression in RA-FLS. RA-FLS and HDMECs co-cultures were respectively treated with As 2 O 3 alone or together with TNF-alpha (100ng/ml) for 48 h. A. TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and HDMECs co-cultures were analyzed by ELISA. Results showed that concentrations of TSP-1, TGF-beta1, CTGF and VEGF increased significantly after treatment with TNF-alpha (100 ng/ml) (n = 3) compared to vehicle control group (n = 3; & p < 0.05), and then significantly decreased concentrations of TSP-1, TGF-beta1, CTGF and VEGF were observed after the treatment of As 2 O 3 at doses of 1.0 uM and 2.0 uM (n = 3, respectively; # p < 0.05, ## p < 0.01). TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and

HDMECs co-cultures without TNF-alpha addition also decreased significantly after treatment of As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) compared to vehicle control (n = 3; * p < 0.05). B. mRNA levels of TSP1, TGF-beta1, CTGF and VEGF expression in RA-FLS co-cultured were performed by real-time PCR. Results showed similar changes of TSP-1, TGF-beta1, CTGF and VEGF mRNA expression as demonstrated in protein regulation after treatment of As 2 O 3 alone (n = 3) or together with TNF-alpha (100ng/ml) (n = 3; * p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01). C. and D. Transwell assay and tube formation test were performed by applying supernatants from RA-FLS and HDMECs co-culture to HDMECs for 6 h respectively. Results showed that migration and capillary-like structure formation of HDMECs significantly increased after TNF-alpha (100ng/ml) stimulation (n = 3, respectively; * p < 0.05), while As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) played a significantly opposite role in migration and tube formation with or without TNF-alpha (* p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01), which were also in a dose dependent manner. E. Ex vivo aortic ring angiogenesis assay showed similar changes of microvessel sprouting as demonstrated in migration and tube formation of HDMECs in the transwell assay and tube formation test above (n = 3, respectively; * p < 0.05, ** p < 0.01, # p < 0.05). Bars = 300um. Original magnification = x5. Results are expressed as the mean ± S.E.M. & p < 0.05 versus Veh. * p < 0.05, ** p < 0.01 versus Veh, # p < 0.05, ## p < 0.01 versus TNF-alpha. Veh = vehicle control under treatment of 1% FBS DMEM alone. TNF-alpha = control group under treatment of TNF-alpha (100ng/ml).Index in PubMed under a CC BY license. PMID: 29088724




A: Effect of AMD3100 on expression of MMP-2, MMP-9 and VEGF in SW480 cells. Protein samples extracted from SW480 cells treated for 26 h with AMD3100 were subjected to Western blotting for MMP-2, MMP-9, VEGF and GAPDH proteins. AMD3100 significantly decreased MMP-9 and VEGF protein expression in SW480 cells in a dose-dependent manner. The level of GAPDH expression was determined as a control for equivalent protein loading; B: RNA samples extracted from SW480 cells treated for 26 h with AMD3100 were subjected to RT-PCR for MMP-2 , MMP-9 , VEGF and beta -actin . AMD3100 also significantly decreased expression of MMP-9 and VEGF mRNAs in SW480 cells, and the inhibitory effect was dose-dependent. RT-PCR for beta -actin was performed in parallel to show an equal amount of total RNA in the sample.Index in PubMed under a CC BY license. PMID: 18416455

As 2 O 3 inhibited angiogenesis by modulating TSP-1, TGF-beta1, CTGF and VEGF expression in RA-FLS. RA-FLS and HDMECs co-cultures were respectively treated with As 2 O 3 alone or together with TNF-alpha (100ng/ml) for 48 h. A. TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and HDMECs co-cultures were analyzed by ELISA. Results showed that concentrations of TSP-1, TGF-beta1, CTGF and VEGF increased significantly




after treatment with TNF-alpha (100 ng/ml) (n = 3) compared to vehicle control group (n = 3; & p < 0.05), and then significantly decreased concentrations of TSP-1, TGF-beta1, CTGF and VEGF were observed after the treatment of As 2 O 3 at doses of 1.0 uM and 2.0 uM (n = 3, respectively; # p < 0.05, ## p < 0.01). TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and HDMECs co-cultures without TNF-alpha addition also decreased significantly after treatment of As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) compared to vehicle control (n = 3; * p < 0.05). B. mRNA levels of TSP1, TGF-beta1, CTGF and VEGF expression in RA-FLS co-cultured were performed by real-time PCR. Results showed similar changes of TSP-1, TGF-beta1, CTGF and VEGF mRNA expression as demonstrated in protein regulation after treatment of As 2 O 3 alone (n = 3) or together with TNF-alpha (100ng/ml) (n = 3; * p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01). C. and D. Transwell assay and tube formation test were performed by applying supernatants from RA-FLS and HDMECs co-culture to HDMECs for 6 h respectively. Results showed that migration and capillary-like structure formation of HDMECs significantly increased after TNF-alpha (100ng/ml) stimulation (n = 3, respectively; * p < 0.05), while As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) played a significantly opposite role in migration and tube formation with or without TNF-alpha (* p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01), which were also in a dose dependent manner. E. Ex vivo aortic ring angiogenesis assay showed similar changes of microvessel sprouting as demonstrated in migration and tube formation of HDMECs in the transwell assay and tube formation test above (n = 3, respectively; * p < 0.05, ** p < 0.01, # p < 0.05). Bars = 300um. Original magnification = x5. Results are expressed as the mean ± S.E.M. & p < 0.05 versus Veh. * p < 0.05, ** p < 0.01 versus Veh, # p < 0.05, ## p < 0.01 versus TNF-alpha. Veh = vehicle control under treatment of 1% FBS DMEM alone. TNF-alpha = control group under treatment of TNF-alpha (100ng/ml).Index in PubMed under a CC BY license. PMID: 29088724




CircPUM1 promotes angiogenesis in ovarian cancer as the form of exosomes. (A) The expression of circPUM1 was positively related with MVD in ovarian cancer tissue. (B) Experimental design schematic (Created in BioRender. Guan, X. (2025) <https://BioRender.com/f5br0q0>): HUVEC were co-cultured with exosomes derived from circPUM1-overexpressing and control OVCAR3 cells. (C) RT-qPCR showed that circPUM1 was highly expressed in HUVEC after co-cultured with exosomal circPUM1. Exosomal circPUM1 enhanced migration ability (D, E), viability (F), invasiveness (G), and tube formation ability (H) of HUVECs. (I) Circular RNA pull-down assays using biotinylated circPUM1 probes confirmed co-enrichment of circPUM1 and miR-607. (J) Dual-luciferase reporter assays showed miR-607 targeting 3'-UTR in both VEGFA and RAB27B. (K) Western blot revealed that overexpression of circPUM1 upregulated VEGFA and RAB27B, whereas miR-607 downregulated these targets in OVCAR3 cells. Exosomal circPUM1 elevated intracellular VEGFA expression in HUVECs. (L) Molecular schematic (Created in BioRender. Guan, X. (2025)


<https://BioRender.com/5eldsz5>): In ovarian cancer, the highly expressed circPUM1 sponges miR-607, upregulating expression of VEGFA and RAB27B. This dual regulation establishes a pro-tumorigenic cascade through two distinct pathways: (1) RAB27B-mediated secretion of exosomal circPUM1, and (2) VEGFA-induced activation of angiogenic signaling. Index in PubMed under a CC BY license. PMID: 41050095




As 2 O 3 inhibited angiogenesis by modulating TSP-1, TGF-beta1, CTGF and VEGF expression in RA-FLS. RA-FLS and HDMECs co-cultures were respectively treated with As 2 O 3 alone or together with TNF-alpha (100ng/ml) for 48 h. A. TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and HDMECs co-cultures were analyzed by ELISA. Results showed that concentrations of TSP-1, TGF-beta1, CTGF and VEGF increased significantly after treatment with TNF-alpha (100 ng/ml) (n = 3) compared to vehicle control group (n = 3; & p < 0.05), and then significantly decreased concentrations of TSP-1, TGF-beta1, CTGF and VEGF were observed after the treatment of As 2 O 3 at doses of 1.0 uM and 2.0 uM (n = 3, respectively; # p < 0.05, ## p < 0.01). TSP-1, TGF-beta1, CTGF and VEGF protein expression in the supernatants from RA-FLS and HDMECs co-cultures without TNF-alpha addition also decreased significantly after treatment of As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) compared to vehicle control (n = 3; * p < 0.05). B. mRNA levels of TSP1, TGF-beta1, CTGF and VEGF expression in RA-FLS co-cultured were performed by real-time PCR. Results showed similar changes of TSP-1, TGF-beta1, CTGF and VEGF mRNA expression as demonstrated in protein regulation after treatment of As 2 O 3 alone (n = 3) or together with TNF-alpha (100ng/ml) (n = 3; * p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01). C. and D. Transwell assay and tube formation test were performed by applying supernatants from RA-FLS and HDMECs co-culture to HDMECs for 6 h respectively. Results showed that migration and capillary-like structure formation of HDMECs significantly increased after TNF-alpha (100ng/ml) stimulation (n = 3, respectively; * p < 0.05), while As 2 O 3 at doses of 1.0 uM (n = 3) and 2.0 uM (n = 3) played a significantly opposite role in migration and tube formation with or without TNF-alpha (* p < 0.05, ** p < 0.01, # p < 0.05, ## p < 0.01), which were also in a dose dependent manner. E. Ex vivo aortic ring angiogenesis assay showed similar changes of microvessel sprouting as demonstrated in migration and tube formation of HDMECs in the transwell assay and tube formation test above (n = 3, respectively; * p < 0.05, ** p < 0.01, # p < 0.05). Bars = 300um. Original magnification = x5. Results are expressed as the mean ± S.E.M. & p < 0.05 versus Veh. * p < 0.05, ** p < 0.01 versus Veh, # p < 0.05, ## p < 0.01 versus TNF-alpha. Veh = vehicle control under treatment of 1% FBS DMEM alone. TNF-alpha = control group under treatment of TNF-alpha (100ng/ml). Index in PubMed under a CC BY license. PMID: 29088724



in HUVECs. (A) QPCR showed a reduction of circPUM1 levels in exosomes derived from P-Nb2C@si-circPUM1-treated ovarian cancer cells. (B) Intracellular circPUM1 level in HUVEC treated with Exo-P-Nb2C@si-circPUM1 were markedly lower than those in Exo-control and Exo-P-Nb2C@si-scramble groups. CCK-8 and EdU assays demonstrated that Exo-P-Nb2C@si-circPUM1 suppressed cell viability (C) and DNA replication capacity (D, E) in HUVECs. Wound healing, Transwell assay and tube formation assay indicated that Exo-P-Nb2C@si-circPUM1 treatment significantly attenuated HUVEC migration (F), invasion (G), and angiogenic potential (I) compared to control groups. Immunofluorescence (H, J) and Western blot (K) confirmed downregulation of VEGFA in Exo-P-Nb2C@si-circPUM1 treated HUVECs. Index in PubMed under a CC BY license. PMID: 41050095



As 2 O 3 suppressed TSP-1, TGF-beta1, CTGF and VEGF expression and microvessel density in synovial tissue of CIA mice. Positive staining appears as brown color. A - D , F. Immunohistochemical analysis demonstrated increased percentage (%) of positive cells for CTGF, TGF-beta1, TSP-1 and VEGF in synovial tissue of CIA mice (n = 6; A-D, II, CIA control mice) compared to synovium of normal mice (n = 6; A-D, I, normal mice; F, *** p < 0.001), while CIA mice treated with As 2 O 3 at a dose of 5.0 mg/kg/day (n = 6; A-D, III) and MTX 1.5mg/kg/week (n = 6; A-D, IV) showed decreased % of positive cells for CTGF, TGF-beta1, TSP-1 and VEGF in synovial tissue compared to CIA control mice (n = 6; A-D, II; F, ### p < 0.001). Original magnification = x40. Bars = 25um. E , G. Immunohistochemical analysis for vWF showed a significant increase in number of microvessels in synovial tissue of CIA control mice (n = 6; E, II) compared to the synovium of normal mice (n = 6; E, I; G, *** p < 0.001), while CIA mice treated with As 2 O 3 at a dose of 5.0 mg/kg/day (n = 6; E, III) and MTX 1.5 mg/kg/week (n = 6; E, IV) demonstrated decreased number of microvessels in synovial tissue compared to CIA control mice (n = 6; E, II; G, ## p < 0.01). Original magnification = x40. Bars = 25um. Data are expressed as the mean ± S.E.M. *** p < 0.001 versus NC, # p < 0.05, ## p < 0.01, ### p < 0.001 versus Con. NC = normal control mice. Con = CIA control mice. MTX = methotrexate. Index in PubMed under a CC BY license. PMID: 29088724



The promotion of NA for the expression of CD31 and VEGF in vivo . (A 1 -A 3) Immunofluorescence staining indicating the CD31 expression in the healed skins on 11 days post-treatment. (A 4) The semi-quantification of CD31 in vivo on day 11. (B 1 -B 3) Immunofluorescence staining indicating the VEGF expression in the healed skins on 11 days post-treatment. (B 4) The semi-quantification of VEGF in vivo on day 11. Statistical significance is indicated as ns p > 0.05, **** p < 0.0001 versus the Control group. Index in PubMed under a CC BY license. PMID: 39502529

108 Publications Citing This Product

dependent gene expression in vivo

2. PubMed ID: -, Sun,S.,Du,Y.,Li,S.,Gao,B.,Xia,R.,Cao,W. ... Zhu,E.(2021).Anti-inflammatory activity of different isolated sites of Chloranthus serratus in complete Freund's adjuvant-induced arthritic rats.Experimental and Therapeutic Medicine,22,848.<https://doi.org/10.3892/etm.2021.10280>

3. PubMed ID: 33785768, Qu X,Fang Y,Zhuang S,Zhang Y.Micro-RNA miR-542-3p suppresses decidualization by targeting ILK pathways in human endometrial stromal cells.Sci Rep.2021 Mar 30;11(1):7186.doi:10.1038/s41598-021-85295-2.PMID:33785768;PMCID:PMC8009905.

Visit bosterbio.com/anti-vegf-antibody-pa1080-boster.html to see all 108 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-VEGF/VEGFA Antibody

For Research Use Only. Not for use in diagnostic procedures.