BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-ERK2/MAPK1 Antibody Picoband®

Catalog Number: PA1310-1

About MAPK1

Mitogen-activated protein kinase 1, also known as MAPK1, p42MAPK, and ERK2, is an enzyme which in humans is encoded by the MAPK1 gene. The gene is mapped to 22q11.2. And The protein encoded by this gene is a member of the MAP kinase family. The deduced 360-amino acid human ERK2 protein shares 98% identity with rat Erk2. The MAP kinase ERK2 is widely involved in eukaryotic signal transduction. Upon activation, it translocates to the nucleus of the stimulated cell, where it phosphorylates nuclear targets. Futhermore, the ERK pathway is necessary for experience-dependent plasticity and for long-term potentiation of synaptic transmission in the developing visual cortex. And the MAPK pathway is also involved in responses to NTN1.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-ERK2/MAPK1 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ERK2/MAPK1 Antibody catalog # PA1310-1. Tested in WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20˚C for one year from date of receipt. After reconstitution, at 4˚C for one month. It can also be aliquotted and stored frozen at -20˚C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P28482

BOSTER
antibody and ELISA experts


BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human ERK2, identical to the related rat and mouse sequences.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Rat, Mouse

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-ERK2/MAPK1 Antibody Picoband® (PA1310-1) Images



Anti-ERK2 antibody, PA1310-1, All Western blottingAll lanes: Anti-MAPK1(PA1310-1) at 0.5ug/ml Lane 1: Human Placenta Tissue Lysate at 40ug Lane 2: Rat Liver Tissue Lysate at 40ug Lane 3: Rat Brain Tissue Lysate at 40ug Lane 4: Rat Thymus Tissue Lysate at 40ug Lane 5: HELA Whole Cell Lysate at 40ug Lane 6: SMMC Whole Cell Lysate at 40ug Lane 7: COLO320 Whole Cell Lysate at 40ug Lane 8: MM231 Whole Cell Lysate at 40ug Lane 9: CEM Whole Cell Lysate at 40ug Predicted bind size: 41KD Observed bind size: 41KD

6 Publications Citing This Product

1. PubMed ID: 26097571, Li Xl, Chen Xq, Zhang Mn, Chen N, Nie L, Xu M, Gong J, Shen Pf, Su Zz, Weng X, Tan Jy, Zhao T, Zeng H, Zhou Q. Int J Clin Exp Pathol. 2015 Apr 1;8(4):3871-81. Ecollection 2015. Sox9 Was Involved In Tkis Resistance In Renal Cell Carcinoma Via Raf/M...

2. PubMed ID: 26097571, Li Xl, Chen Xq, Zhang Mn, Chen N, Nie L, Xu M, Gong J, Shen Pf, Su Zz, Weng X, Tan Jy, Zhao T, Zeng H, Zhou Q. Int J Clin Exp Pathol. 2015 Apr 1;8(4):3871-81. Ecollection 2015. Sox9 Was Involved In Tkis Resistance In Renal Cell Carcinoma Via Raf/M...

3. PubMed ID: 21769431, Sang J, Chen Y, Tao Y. Mol Med Rep. 2011 Nov-Dec;4(6):1163-7. Doi: 10.3892/Mmr.2011.535. Epub 2011 Jul 15. Nitric Oxide Inhibits Gastric Cancer Cell Growth Through The Modulation Of The Akt Pathway.

Visit bosterbio.com/anti-erk2-antibody-pa1310-1-boster.html to see all 6 publications.

For Research Use Only. Not for use in diagnostic procedures.