

Anti-MAPK8/9 Antibody

Catalog Number: PA1407

About MAPK8

Mitogen-activated protein kinase 8 is an enzyme that in humans is encoded by the MAPK8 gene. The protein encoded by this gene is a member of the MAP kinase family. The International Radiation Hybrid Mapping Consortium mapped the MAPK8 gene to chromosome 10. The single long open reading frame encodes a putative protein kinase with a predicted mass of 44.2 kD. This kinase play a key role in T cell proliferation, apoptosis and differentiation. The protein encoded by MAPK9 gene is also a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase targets specific transcription factors, and thus mediates immediate-early gene expression in response to various cell stimuli. It is most closely related to MAPK8, both of which are involved in UV radiation induced apoptosis, thought to be related to the cytochrome c-mediated cell death pathway. This gene and MAPK8 are also known as c-Jun N-terminal kinases. This kinase blocks the ubiquitination of tumor suppressor p53, and thus it increases the stability of p53 in nonstressed cells. Studies of this gene's mouse counterpart suggest a key role in T-cell differentiation. Several alternatively spliced transcript variants encoding distinct isoforms have been reported.

Overview

Product Name	Anti-MAPK8/9 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-MAPK8/9 Antibody catalog # PA1407. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P45983, P45984

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human MAPK8/9, identical to the related mouse and rat sequences.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).






888-466-3604 | support@bosterbio.com | www.bosterbio.com

Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Rat, Mouse, By Heat - Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat




Anti-MAPK8/9 Antibody (PA1407) Images




Anti-MAPK8/9 antibody, PA1407, Western blotting

Lane 1: Rat Brain Tissue Lysate Lane 2: Rat Thymus Tissue Lysate

Lane 3: MCF-7 Cell Lysate Lane 4: HELA Cell Lysate Lane 5: JURKAT Cell Lysate Lane 6: MM231 Cell Lysate Lane 7: CEM Cell Lysate



Anti-MAPK8/9 antibody, PA1407, IHC(P) IHC(P): Rat Intestine Tissue



Anti-MAPK8/9 antibody, PA1407, IHC(P) IHC(P): Human Intestinal Cancer Tissue

6 Publications Citing This Product

1. PubMed ID: 21625440, Ding Y, Zou J, Li Z, Tian J, Abdelalim S, Du F, She R, Wang D, Tan C, Wang H, Chen W, Lv D, Chang L. Plos One. 2011;6(5):E20008. Doi: 10.1371/Journal.Pone.0020008. Epub 2011 May 23. Study Of Histopathological And Molecular Changes Of Rat Kidney Un...

- 2. PubMed ID: 29072695, Glucocorticoid mediates prenatal caffeine exposure-induced endochondral ossification retardation and its molecular mechanism in female fetal rats
- 3. PubMed ID: 29399151, Pseudolaric acid B induced autophagy, but not apoptosis, in MRC5 human fibroblast cells


Visit bosterbio.com/anti-mapk8-9-antibody-pa1407-boster.html to see all 6 publications.

Submit a product review to Biocompare.com











reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-MAPK8/9 Antibody